

ON THE MOTION OF FREE INTERFACE IN TWO-DIMENSIONAL IDEAL INCOMPRESSIBLE MHD

CHENGCHUN HAO

ABSTRACT. For the free boundary problem of the plasma-vacuum interface to ideal incompressible magnetohydrodynamics (MHD) in two-dimensional space, the a priori estimates of solutions are proved in Sobolev norms by adopting a geometrical point of view. In the vacuum region, the magnetic field is described by the div-curl system of pre-Maxwell dynamics, while at the interface the total pressure is continuous and the magnetic field is tangent to the boundary. We prove that the L^2 norms of any order covariant derivatives of the magnetic field in vacuum and on the boundaries are bounded in terms of initial data and the second fundamental forms of the free interface and the rigid wall.

1. INTRODUCTION

In the present paper, we consider the free boundary problem of 2D ideal incompressible magnetohydrodynamics (MHD). It consists of finding a bounded variable domain $\Omega_t^+ \subset \mathbb{R}^2$ filled with inviscid incompressible electrically conducting homogeneous plasma (the density $\rho(t, x) \equiv \text{const} > 0$), together with the vector field of velocity $\mathbf{v}(t, x) = (v_1, v_2)$, the scalar pressure $p(t, x)$ and the magnetic field $\mathbf{H}(t, x) = (H_1, H_2)$ satisfying the system of MHD. The boundary Γ_t of Ω_t^+ is the free surface of the plasma. It is assumed that the plasma is surrounded by a vacuum region Ω_t^- and that the whole domain $\Omega = \Omega_t^+ \cup \Gamma_t \cup \Omega_t^-$ is independent of time and bounded by a fixed perfectly conducting rigid wall W such that $W \cap \Gamma_0 = \emptyset$. Both Ω_t^+ and Ω are simply connected, while Ω^- is not simply connected in 2D rather than 3D. The magnetic field should be found not only in Ω_t^+ but also in Ω_t^- .

In the plasma region Ω_t^+ , the ideal MHD equations apply, i.e., for $t > 0$

$$\begin{cases} \mathbf{v}_t + \mathbf{v} \cdot \nabla \mathbf{v} + \nabla p = \mu(\mathbf{H} \cdot \nabla \mathbf{H} - \frac{1}{2} \nabla |\mathbf{H}|^2), & (1.1a) \\ \mathbf{H}_t + \mathbf{v} \cdot \nabla \mathbf{H} = \mathbf{H} \cdot \nabla \mathbf{v}, & (1.1b) \\ \text{div } \mathbf{v} = 0, \quad \text{div } \mathbf{H} = 0. & (1.1c) \end{cases}$$

Let $\hat{\mathbf{H}} = (\hat{H}_1, \hat{H}_2)$ be the magnetic field in the vacuum Ω_t^- . Since the vacuum has no density, velocity, electric current (i.e., $\hat{E}_t = 0$), except for the magnetic field, we have the pre-Maxwell equations in vacuum

$$\partial_1 \hat{H}_2 = \partial_2 \hat{H}_1, \quad \text{div } \hat{\mathbf{H}} = 0, \quad \hat{\mathbf{H}}_t = -\nabla^\perp \hat{E},$$

where $\nabla^\perp = (\partial_2, -\partial_1)$, and the electric field $\hat{E}$ is scalar, a secondary variable, which is different from the 3D case. At the wall W , the tangential component of the electric field and the normal component of the magnetic field must vanish

$$\hat{E} = 0, \quad \mathbf{n} \cdot \hat{\mathbf{H}} = 0, \quad \text{on } W,$$

where $\mathbf{n}$ is the inward drawn unit normal to the boundary of Ω_t^- .

The plasma-vacuum interface is now free to move since the plasma is surrounded by vacuum. Hence, $\mathbf{v} \cdot \mathbf{n}|_{\Gamma_t}$ is unknown and arbitrary where $\mathbf{n}$ is the unit normal to Γ_t pointing from the plasma to the vacuum. Thus, we need some non-trivial jump conditions that

must be satisfied to connect the fields across the interface. These arise from the divergence $\mathbf{H}$ equation, Faraday's law and the momentum equation. A convenient way to obtain the desired relations is to assume that the plasma surface Γ_t is moving with a normal velocity

$$V_{\mathbf{n}}\mathbf{n} = (\mathbf{v} \cdot \mathbf{n})\mathbf{n},$$

where $V_{\mathbf{n}}$ is the velocity of evolution of Γ_t in the direction $\mathbf{n}$. The jump conditions are straightforward to derive in a reference frame moving with the fluid surface. Once these conditions are obtained, all that is then required is to convert back to the laboratory frame using the corresponding Galilean transformation (cf. [7]). From Maxwell's equations, we know that, at the interface Γ_t , the magnetic field and the electric field must satisfy the conditions

$$\mathbf{n} \cdot \mathbf{H} = \mathbf{n} \cdot \hat{\mathbf{H}} = 0, \quad \hat{E}\mathbf{n}^\perp = (\mathbf{v} \cdot \mathbf{n})\mathbf{H}, \quad \text{on } \Gamma_t, \quad (1.2)$$

where $\mathbf{n}^\perp = (n_2, -n_1)$. The first one also means that the magnetic fields are not pointing into the vacuum on the interface.

We also have the following pressure balance condition (also cf. [7, 17]) on the interface Γ_t :

$$p + \frac{\mu}{2}|\mathbf{H}|^2 = \frac{\mu}{2}|\hat{\mathbf{H}}|^2, \quad \text{on } \Gamma_t.$$

For convenience, we denote

$$P = q^+ - q^-, \quad q^+ = p + \frac{\mu}{2}|\mathbf{H}|^2, \quad q^- = \frac{\mu}{2}|\hat{\mathbf{H}}|^2.$$

The system can be rewritten as

$$\begin{cases} \mathbf{v}_t + \mathbf{v} \cdot \nabla \mathbf{v} + \nabla q^+ = \mu \mathbf{H} \cdot \nabla \mathbf{H}, & x \in \Omega_t^+, & (1.3a) \\ \mathbf{H}_t + \mathbf{v} \cdot \nabla \mathbf{H} = \mathbf{H} \cdot \nabla \mathbf{v}, & x \in \Omega_t^+, & (1.3b) \\ \operatorname{div} \mathbf{v} = 0, \quad \operatorname{div} \mathbf{H} = 0, & x \in \Omega_t^+, & (1.3c) \\ \nabla^\perp \cdot \hat{\mathbf{H}} = 0, \quad \operatorname{div} \hat{\mathbf{H}} = 0, \quad \hat{\mathbf{H}}_t = -\nabla^\perp \hat{E}, & x \in \Omega_t^-, & (1.3d) \\ P = 0, \quad \mathbf{H} \cdot \mathbf{n} = \hat{\mathbf{H}} \cdot \mathbf{n} = 0, & x \in \Gamma_t, & (1.3e) \\ \hat{E}\mathbf{n}^\perp = (\mathbf{v} \cdot \mathbf{n})\hat{\mathbf{H}}, & x \in \Gamma_t, & (1.3f) \\ \hat{E} = 0, \quad \hat{\mathbf{H}} \cdot \mathbf{n} = 0, & x \in W, & (1.3g) \\ \mathbf{v}(0, x) = \mathbf{v}_0(x), \quad \mathbf{H}(0, x) = \mathbf{H}_0(x), & x \in \Omega^+, & (1.3h) \\ \hat{\mathbf{H}}(0, x) = \hat{\mathbf{H}}_0(x), \quad \hat{E}(0, x) = \hat{E}_0(x), & x \in \Omega^-, & (1.3i) \\ \Omega_t^+|_{t=0} = \Omega^+, \quad \Omega_t^-|_{t=0} = \Omega^-, \quad \Gamma_t|_{t=0} = \Gamma. & & (1.3j) \end{cases}$$

We will prove a priori bounds for the interface problem (1.3) in Sobolev spaces under the following condition

$$\nabla_{\mathbf{n}}P \leq -\varepsilon < 0 \text{ on } \Gamma, \quad (1.4)$$

where $\nabla_{\mathbf{n}} = \mathbf{n}^i \partial_i$. In fact, we assume that this condition holds initially, and then we can verify that it holds true within a period. For the free boundary problem of motion of incompressible fluids in vacuum, without magnetic fields, the natural physical condition (cf. [2, 4, 5, 12, 13, 14, 6, 20, 24, 25, 26]) reads that $\nabla_{\mathbf{n}}p \leq -\varepsilon < 0$ on Γ , which excludes the possibility of the Rayleigh-Taylor type instability (see [6]).

There were few well-posedness results for full non-stationary plasma-vacuum models because of the difficulties caused by the strong coupling between the velocity field and the magnetic field. Hao and Luo studied the a priori estimates for the free boundary problem of ideal incompressible MHD flows in [9] with a bounded initial domain homeomorphic to a ball, provided that the size of the magnetic field to be invariant on the free boundary

that includes the case of zero magnetic boundary condition. For the special case when the magnetic field is zero on the free boundary, Lee proved the local existence and uniqueness of plasma-vacuum free boundary problem of incompressible viscous-diffusive MHD flow in three-dimensional space with infinite depth setting in [10], and got a local unique solution of free boundary MHD without kinetic viscosity and magnetic diffusivity via zero kinetic viscosity-magnetic diffusivity limit in [11]. For the incompressible viscous MHD equations, a free boundary problem in a simply connected domain of $\mathbb{R}^3$ was studied by a linearization technique and the construction of a sequence of successive approximations in [16] with an irrotational condition for magnetic fields in a part of the domain. The well-posedness of the linearized plasma-vacuum interface problem in ideal incompressible MHD was studied in [15] in an unbounded plasma domain, the linearized plasma-vacuum problem in ideal compressible MHD was investigated in [23, 18], the well-posedness of the original nonlinear free boundary problem was proved in [19] by using the Nash-Moser method, and a stationary problem was studied in [8]. The current-vortex sheets problem was studied in [3], [21] and [22].

Regarding the cases without magnetic fields, the free surface problem of the incompressible Euler equations of fluids has attracted much attention in recent decades and important progress has been made for flows with or without vorticity or surface tension. We refer readers to [1, 4, 5, 12, 13, 14, 6, 20, 24, 25, 26] and references therein.

In this paper, we prove a priori bounds for the free interface problem (1.3) and take into account the magnetic field not only in plasma but also in vacuum. Thus, we do not need the restricted boundary condition $|\mathbf{H}| \equiv \text{const} \geq 0$ on the free interface assumed in [9]. What makes this problem difficult is that the regularity of the boundary enters to the highest order and energies exchange between plasma and vacuum. Roughly speaking, the energies of plasma and vacuum will exchange via the pressure balance relation $q^+ = q^-$ on the free interface. But we need to investigate the estimates of the magnetic field and the electric field in vacuum even if the electric field is a secondary variable in order to obtain energy estimates. We also introduce the “virtual particle” endowed with a virtual velocity field to reformulate the free boundary problem to a fixed boundary problem. We can show that the norms of the magnetic field in vacuum depend only on the norms of initial data and the second fundamental forms of the interface Γ_t and the wall W .

Now, we derive the conserved energy. Let $D_t := \partial_t + v^k \partial_k$ be the material derivative, then it holds for any function F on $\overline{\Omega_t^+}$

$$\frac{d}{dt} \int_{\Omega_t^+} F dx = \int_{\Omega_t^+} D_t F dx, \quad (1.5)$$

since $\text{div } \mathbf{v} = 0$, and then for any function F on $\overline{\Omega_t^-}$ (we can extend it to Ω by a smooth cut-off function)

$$\frac{d}{dt} \int_{\Omega_t^-} F dx = \int_{\Omega_t^-} \partial_t F dx - \int_{\Gamma_t} \mathbf{v}_n F dS, \quad (1.6)$$

where $\mathbf{n}$ is the outward unit normal to Γ_t corresponding to Ω_t^+ .

Throughout the paper, we use the Einstein summation convention, that is, when an index variable appears twice in both the subscript and the superscript of a single term it indicates summation of that term over all the values of the index.

From (1.5) and (1.6), by using the boundary conditions (1.3e), (1.3f) and (1.3g), the conserved physical energy is given by

$$E_0(t) := \int_{\Omega_t^+} \left(\frac{1}{2} |\mathbf{v}(t)|^2 + \frac{\mu}{2} |\mathbf{H}(t)|^2 \right) dx + \int_{\Omega_t^-} \frac{\mu}{2} |\hat{\mathbf{H}}(t)|^2 dx \equiv E_0(0),$$

for any $t > 0$.

The higher order energy has an interface boundary part and interior parts in both plasma and vacuum. The boundary part controls the norms of the second fundamental form of the free interface, the interior part in plasma controls the norms of the velocity, magnetic fields and hence the pressure, the interior part in vacuum controls the magnetic field. We will prove that the time derivatives of the energy are controlled by themselves. A crucial point in the construction of the higher order energy norms is that the time derivatives of the interior parts will, after integrating by parts, contribute some boundary terms that cancel the leading-order terms in the corresponding time derivatives of the boundary integrals. To this end, we need to project the equations for the total pressure P to the tangent space of the boundary.

The higher order energies are defined as: For $r \geq 1$

$$E_r(t) = \int_{\Omega_t^+} \delta^{ij} Q(\partial^r v_i, \partial^r v_j) dx + \mu \int_{\Omega_t^+} \delta^{ij} Q(\partial^r H_i, \partial^r H_j) dx \\ + \operatorname{sgn}(r-1) \int_{\Gamma_t} Q(\partial^r P, \partial^r P) \vartheta dS + \int_{\Omega_t^+} (|\partial^{r-1}(\nabla^\perp \cdot v)|^2 + \mu |\partial^{r-1}(\nabla^\perp \cdot B)|^2) dx,$$

where $\operatorname{sgn}(s)$ be the sign function of the real number s and $\vartheta = (-\nabla_{\mathbf{n}} P)^{-1}$. Here, Q is a positive definite quadratic form which, when restricted to the boundary, is the inner product of the tangential components, i.e., $Q(\alpha, \beta) = \langle \Pi \alpha, \Pi \beta \rangle$, and in the interior $Q(\alpha, \alpha)$ increases to the norm $|\alpha|^2$. To be more specific, let

$$Q(\alpha, \beta) = q^{i_1 j_1} \dots q^{i_r j_r} \alpha_{i_1 \dots i_r} \beta_{j_1 \dots j_r}$$

where

$$q^{ij} = \delta^{ij} - \eta(d)^2 \mathbf{n}^i \mathbf{n}^j, \quad d(x) = \operatorname{dist}(x, \Gamma_t), \quad \mathbf{n}^i = -\delta^{ij} \partial_j d.$$

Here η is a smooth cutoff function satisfying $0 \leq \eta(d) \leq 1$, $\eta(d) = 1$ when $d < d_0/4$ and $\eta(d) = 0$ when $d > d_0/2$. d_0 is a fixed number that is smaller than the injectivity radius of the normal exponential map ι_0 , defined to be the largest number ι_0 such that the map

$$\Gamma_t \times (-\iota_0, \iota_0) \rightarrow \{x \in \mathbb{R}^2 : \operatorname{dist}(x, \Gamma_t) < \iota_0\}, \quad (1.7)$$

given by

$$(\bar{x}, \iota) \rightarrow x = \bar{x} + \iota \mathbf{n}(\bar{x}),$$

is an injection. We can also define ι_0 for W similarly which is independent of t and denote it by ι'_0 . We always assume the second fundamental form of W and $1/\iota'_0$ to be bounded, since both of them are invariant with respect to time.

Now, we can state the main theorem as follows.

Theorem 1.1. *Let*

$$\mathcal{K}(0) = \max(\|\theta(0, \cdot)\|_{L^\infty(\Gamma)}, 1/\iota_0(0)), \quad (1.8)$$

$$\mathcal{E}(0) = \|1/(\nabla_N P(0, \cdot))\|_{L^\infty(\Gamma)} = 1/\varepsilon > 0. \quad (1.9)$$

Assume $\|\theta\|_{L^\infty(W)} + 1/\iota'_0 \leq C\mathcal{K}(0)$, $\|\nabla_N P(0, \cdot)\|_{L^\infty(\Gamma)} \leq C$ and $\|\nabla^\perp \hat{E}_0\|_{L^2(\Omega^-)} \leq C$. Then there exists a continuous function $\mathcal{T} > 0$ such that if

$$T \leq \mathcal{T}(\mathcal{K}(0), \mathcal{E}(0), E_0(0), \dots, E_3(0), \operatorname{Vol} \Omega), \quad (1.10)$$

then any smooth solution of the free boundary problem for 2D MHD equations (1.3) with (1.4) in $[0, T]$ satisfies the estimate

$$\sum_{s=0}^3 E_s(t) \leq 2 \sum_{s=0}^3 E_s(0), \quad 0 \leq t \leq T. \quad (1.11)$$

The rest of this paper is organized as follows: In Section 2, we use the Lagrangian coordinates to transform the free interface problem to a fixed initial boundary problem. Sections 3 and 4 are devoted to the estimates of the magnetic field and the electric field in vacuum, respectively. In Sections 5–7, we prove the higher orders energy estimates. Some a priori assumptions will be justified in Section 8. In order to make this paper more readable, we give an appendix on some estimates used in previous sections, which are basically proved in [4].

2. REFORMULATION IN LAGRANGIAN COORDINATES

We may think that the velocity field of the “virtual particles” in vacuum is $\mathbf{v}$ on the boundary. Then, we can extend the velocity from the boundary to the interior of vacuum by a cut-off function such that

$$\mathbf{v}(t, x) = \begin{cases} \mathbf{v}(t, \bar{x}), & \text{near } \Gamma_t, \\ \text{smooth}, & \text{otherwise,} \\ 0, & \text{near } W, \end{cases}$$

and $\operatorname{div} \mathbf{v} = 0$ for $x \in \Omega_t^-$ as long as $\Gamma_t \cap W = \emptyset$ in $[0, T]$, where $\bar{x} \in \Gamma_t$ satisfies $\operatorname{dist}(x, \Gamma_t) = |x - \bar{x}|$, and $|\mathbf{v}(t, x)| \leq \|\mathbf{v}(t)\|_{L^\infty(\Gamma_t)}$ for $x \in \Omega_t^-$ by construction.

Assume that we are given a velocity vector field $\mathbf{v}(t, x)$ defined in a set $\mathcal{D} \subset [0, T] \times \mathbb{R}^n$ such that the interface of $\Omega_t^+ = \{x : (t, x) \in \mathcal{D}\}$ and Ω_t^- moves with the velocity, i.e., $(1, \mathbf{v}) \in T(\partial\mathcal{D})$ which denotes the tangent space of $\partial\mathcal{D}$. We will now introduce Lagrangian or co-moving coordinates, that is, coordinates that are constant along the integral curves of the velocity vector field so that the boundary becomes fixed in these coordinates (cf. [4]). Let $x = x(t, y) = f_t(y)$ be the trajectory of the particles given by

$$\begin{cases} \frac{dx}{dt} = \mathbf{v}(t, x(t, y)), & (t, y) \in [0, T] \times \Omega, \\ x(0, y) = f_0(y), & y \in \Omega. \end{cases} \quad (2.1)$$

where, when $t = 0$, we can start with either the Euclidean coordinates in Ω or some other coordinates $f_0 : \Omega \rightarrow \Omega$ where f_0 is a diffeomorphism in which the domain Ω becomes simple. For simplicity, we will assume $f_0(y) = y$ in this paper. Then, the Jacobian determinant $\det(\partial y / \partial x) \equiv 1$ due to the divergence-free property of $\mathbf{v}$. For each t , we will then have a change of coordinates $f_t : \Omega \rightarrow \Omega$, taking $y \rightarrow x(t, y)$. The Euclidean metric δ_{ij} in Ω then induces a metric

$$g_{ab}(t, y) = \delta_{ij} \frac{\partial x^i}{\partial y^a} \frac{\partial x^j}{\partial y^b} \quad (2.2)$$

in Ω for each fixed t . We use the covariant differentiation in Ω with respect to the metric $g_{ab}(t, y)$, because it corresponds to differentiation in Ω under the change of coordinates $\Omega \ni y \rightarrow x(t, y) \in \Omega$, and we will work in both coordinate systems. This also avoids possible singularities in the change of coordinates.

We recall a result concerning time derivatives of the change of coordinates and commutators between time derivatives and space derivatives (cf. [4, 9, Lemma 2.1]).

Lemma 2.1. *Let $x = f_t(y)$ be the change of variables given by (2.1), and let g_{ab} be the metric given by (2.2). Let $v_i = \delta_{ij} v^j = v^i$, and set*

$$u_a(t, y) = v_i(t, x) \frac{\partial x^i}{\partial y^a}, \quad u^a = g^{ab} u_b, \quad (2.3)$$

$$h_{ab} = \frac{1}{2} D_t g_{ab}, \quad h^{ab} = g^{ac} h_{cd} g^{db}. \quad (2.4)$$

Then

$$D_t \frac{\partial x^i}{\partial y^a} = \frac{\partial x^k}{\partial y^a} \frac{\partial v^i}{\partial x^k}, \quad D_t \frac{\partial y^a}{\partial x^i} = -\frac{\partial y^a}{\partial x^k} \frac{\partial v^k}{\partial x^i}, \quad (2.5)$$

$$D_t g_{ab} = \nabla_a u_b + \nabla_b u_a, \quad D_t g^{ab} = -2h^{ab}, \quad D_t d\mu_g = \text{tr } h d\mu_g, \quad (2.6)$$

where $d\mu_g$ is the Riemannian volume element on Ω in the metric g .

We now recall the estimates of commutators between the material derivative D_t and covariant derivatives ∇_a .

Lemma 2.2 (cf. [4, 9]). *Let $T_{a_1 \dots a_r}$ be a $(0, r)$ tensor. We have*

$$[D_t, \nabla_a] T_{a_1 \dots a_r} = -(\nabla_{a_1} \nabla_a u^d) T_{da_2 \dots a_r} - \dots - (\nabla_{a_r} \nabla_a u^d) T_{a_1 \dots a_{r-1} d}. \quad (2.7)$$

If $\Delta = g^{cd} \nabla_c \nabla_d$ and q is a function, we have

$$[D_t, g^{ab} \nabla_a] T_b = -2h^{ab} \nabla_a T_b - (\Delta u^e) T_e, \quad (2.8)$$

$$[D_t, \nabla] q = 0, \quad (2.9)$$

$$[D_t, \Delta] q = -2h^{ab} \nabla_a \nabla_b q - (\Delta u^e) \nabla_e q. \quad (2.10)$$

Furthermore, for $r \geq 2$,

$$[D_t, \nabla^r] q = \sum_{s=1}^{r-1} -\mathfrak{L}_r^{s+1} (\nabla^{s+1} u) \cdot \nabla^{r-s} q, \quad (2.11)$$

where the symmetric dot product is defined to be in components

$$((\nabla^{s+1} u) \cdot \nabla^{r-s} q)_{a_1 \dots a_r} = \frac{1}{r!} \sum_{\sigma \in \Sigma_r} \left(\nabla_{a_{\sigma_1} \dots a_{\sigma_{s+1}}}^{s+1} u^d \right) \nabla_{da_{\sigma_{s+2}} \dots a_{\sigma_r}}^{r-s} q.$$

Denote

$$\begin{aligned} H_i &= \delta_{ij} H^j = H^i, \quad \beta_a = H_j \frac{\partial x^j}{\partial y^a}, \quad \beta^a = g^{ab} \beta_b, \quad |\beta|^2 = \beta_a \beta^a, \\ \hat{H}_i &= \delta_{ij} \hat{H}^j = \hat{H}^i, \quad \varpi_a = \hat{H}_j \frac{\partial x^j}{\partial y^a}, \quad \varpi^a = g^{ab} \varpi_b, \quad |\varpi|^2 = \varpi_a \varpi^a. \end{aligned}$$

It follows that

$$|\beta| = |\mathbf{H}|, \quad |\varpi| = |\hat{\mathbf{H}}|, \quad H_j = \frac{\partial y^a}{\partial x^j} \beta_a, \quad \hat{H}_j = \frac{\partial y^a}{\partial x^j} \varpi_a. \quad (2.12)$$

From (2.5), we have

$$\begin{aligned} D_t \varpi_a &= D_t \left(\hat{H}_j \frac{\partial x^j}{\partial y^a} \right) = \frac{\partial x^j}{\partial y^a} D_t \hat{H}_j + \hat{H}_j D_t \frac{\partial x^j}{\partial y^a} \\ &= \frac{\partial x^j}{\partial y^a} \left(-(\nabla^\perp \hat{E})_j + v^k \partial_k \hat{H}_j \right) + \hat{H}_j \frac{\partial x^k}{\partial y^a} \frac{\partial v^j}{\partial x^k} \\ &= \frac{\partial x^j}{\partial y^a} (-\nabla^\perp \hat{E})_j + \frac{\partial x^j}{\partial y^a} \frac{\partial x^k}{\partial y^b} \frac{\partial y^b}{\partial x^l} v^l \partial_k \hat{H}_j + \hat{H}_j \frac{\partial x^k}{\partial y^a} \frac{\partial v^l}{\partial x^k} \frac{\partial x^j}{\partial y^b} \frac{\partial y^b}{\partial x^l} \\ &= -\frac{\partial x^j}{\partial y^a} (\nabla^\perp \hat{E})_j + u^b \nabla_b \varpi_a + \varpi_b \nabla_a u^b. \end{aligned}$$

Due to $\det(\partial y / \partial x) \equiv 1$, we get

$$\frac{\partial x^j}{\partial y^a} (\nabla^\perp \hat{E})_j = -\frac{\partial x^1}{\partial y^a} \frac{\partial \hat{E}}{\partial x^2} + \frac{\partial x^2}{\partial y^a} \frac{\partial \hat{E}}{\partial x^1} = \nabla_a^\perp \Xi,$$

where $\nabla^\perp = (\nabla_2, -\nabla_1)$ in Lagrangian coordinates.

Thus, we have obtained

$$D_t \varpi_a = -\nabla_a^\perp \Xi + u^b \nabla_b \varpi_a + \varpi_b \nabla_a u^b.$$

Similarly, we get

$$\nabla_1 \varpi_2 = \nabla_2 \varpi_1.$$

We also have those equations for $D_t u_a$ and $D_t \beta_a$, one can see [9] for details.

Thus, the system (1.3) can be written in the Lagrangian coordinates, for $t > 0$, as

$$\begin{cases} D_t u_a + \nabla_a q^+ = u^c \nabla_a u_c + \mu \beta^d \nabla_d \beta_a, & \text{in } \Omega^+, & (2.13a) \\ D_t \beta_a = \beta^d \nabla_d u_a + \beta^c \nabla_a u_c, & \text{in } \Omega^+, & (2.13b) \\ D_t \varpi_a = -\nabla_a^\perp \Xi + u^b \nabla_b \varpi_a + \varpi_b \nabla_a u^b, & \text{in } \Omega^-, & (2.13c) \\ \nabla_a u^a = 0 \text{ and } \nabla_a \beta^a = 0, & \text{in } \Omega^+, & (2.13d) \\ \nabla_a u^a = 0, \nabla_1 \varpi_2 = \nabla_2 \varpi_1, \nabla_a \varpi^a = 0, & \text{in } \Omega^-, & (2.13e) \\ P = 0, \quad \beta_a N^a = \varpi_a N^a = 0, & \text{on } \Gamma, & (2.13f) \\ \Xi N^\perp = u_N \varpi, & \text{on } \Gamma, & (2.13g) \\ u = 0, \varpi_a N^a = 0, \Xi = 0, & \text{on } W, & (2.13h) \end{cases}$$

where N is the unit normal vector pointing into the interior of Ω^- .

Finally, the energy defined by

$$E_0(t) = \int_{\Omega^+} \left(\frac{1}{2} |u(t)|^2 + \frac{\mu}{2} |\beta(t)|^2 \right) d\mu_g + \int_{\Omega^-} \frac{\mu}{2} |\varpi(t)|^2 d\mu_g$$

is conserved. Of course, it is the equivalent one as in Eulerian coordinates. It can be easily verified by using the Gauss formula:

$$\int_{\Omega^+} \nabla_a F^a d\mu_g = \int_{\Gamma} N_a F^a d\mu_\gamma, \quad \int_{\Omega^-} \nabla_a F^a d\mu_g = - \int_{\Gamma \cup W} N_a F^a d\mu_\gamma, \quad (2.14)$$

where F is a smooth vector-valued function, and (N_a) denotes the outward (or inward) unit conormal to Γ (or W) corresponding to Ω^+ (or Ω^-).

3. THE ESTIMATES OF THE MAGNETIC FIELD IN VACUUM

Since $\nabla_a \varpi^a = 0$ and $\nabla_1 \varpi_2 = \nabla_2 \varpi_1$, we have the following L^2 -estimates.

Proposition 3.1. *Let $r \geq 0$ be an integer. If $|\theta| + 1/\iota_0 \leq K$ on $\Gamma \cup W$, $|\nabla N| \leq CK$ in Ω^- , then it holds*

$$\|\nabla^{r+1} \varpi\|_{L^2(\Omega^-)}^2 \leq CK^{2(r+1)} E_0(0), \quad \|\nabla^r \varpi\|_{L^2(\Gamma \cup W)}^2 \leq CK^{2r+1} E_0(0).$$

Proof. By the Hölder inequality, we have

$$\begin{aligned} \|\nabla \varpi\|_{L^2(\Omega^-)}^2 &= \int_{\Omega^-} g^{ab} g^{cd} \nabla_a \varpi_c \nabla_d \varpi_b d\mu_g \\ &= \int_{\Omega^-} g^{ab} g^{cd} \nabla_a (\varpi_c \nabla_d \varpi_b) d\mu_g - \int_{\Omega^-} g^{cd} \varpi_c \Delta \varpi_d d\mu_g \\ &= - \int_{\Gamma \cup W} N^b g^{cd} \varpi_c \nabla_d \varpi_b d\mu_\gamma \\ &= - \int_{\Gamma \cup W} g^{cd} \varpi_c \nabla_d (N^b \varpi_b) d\mu_\gamma + \int_{\Gamma \cup W} g^{cd} \varpi_c \nabla_d N^b \varpi_b d\mu_\gamma \\ &= - \int_{\Gamma \cup W} \varpi_N \nabla_N \varpi_N d\mu_\gamma - \int_{\Gamma \cup W} \gamma^{cd} \varpi_c \nabla_d \varpi_N d\mu_\gamma + \int_{\Gamma \cup W} g^{cd} \varpi_c \nabla_d N^b \varpi_b d\mu_\gamma \end{aligned}$$

$$\begin{aligned}
&= \int_{\Gamma \cup W} \gamma^{cd} \varpi_c \nabla_d N^b \varpi_b d\mu_\gamma \\
&\leq C \|\theta\|_{L^\infty(\Gamma \cup W)} \|\varpi\|_{L^2(\Gamma \cup W)}^2.
\end{aligned}$$

We get, from Gauss' formula, Hölder's inequalities and Cauchy's inequality, that

$$\begin{aligned}
\|\varpi\|_{L^2(\Gamma \cup W)}^2 &= \int_{\Gamma \cup W} \varpi^c \varpi_c d\mu_\gamma = \int_{\Gamma \cup W} N_a N^a \varpi^c \varpi_c d\mu_\gamma \\
&= - \int_{\Omega^-} \nabla_a (N^a \varpi^c \varpi_c) d\mu_g \\
&= - \int_{\Omega^-} \nabla_a N^a \varpi^c \varpi_c d\mu_g - 2 \int_{\Omega^-} N^a \nabla_a \varpi^c \varpi_c d\mu_g \\
&\leq C \|\operatorname{tr}(\nabla N)\|_{L^\infty(\Omega^-)} \|\varpi\|_{L^2(\Omega^-)}^2 + C \|\nabla \varpi\|_{L^2(\Omega^-)} \|\varpi\|_{L^2(\Omega^-)} \quad (3.1) \\
&\leq C \|\operatorname{tr}(\nabla N)\|_{L^\infty(\Omega^-)} \|\varpi\|_{L^2(\Omega^-)}^2 + \delta \|\nabla \varpi\|_{L^2(\Omega^-)}^2 + C/\delta \|\varpi\|_{L^2(\Omega^-)}^2,
\end{aligned}$$

for any $\delta > 0$. Thus, it follows that

$$\begin{aligned}
\|\nabla \varpi\|_{L^2(\Omega^-)}^2 &\leq C \|\theta\|_{L^\infty(\Gamma \cup W)} \|\operatorname{tr}(\nabla N)\|_{L^\infty(\Omega^-)} \|\varpi\|_{L^2(\Omega^-)}^2 + C\delta \|\theta\|_{L^\infty(\Gamma \cup W)} \|\nabla \varpi\|_{L^2(\Omega^-)}^2 \\
&\quad + C/\delta \|\theta\|_{L^\infty(\Gamma \cup W)} \|\varpi\|_{L^2(\Omega^-)}^2.
\end{aligned}$$

Taking δ so small that $C\delta K = 1/2$, we get

$$\begin{aligned}
\|\nabla \varpi\|_{L^2(\Omega^-)}^2 &\leq C \|\theta\|_{L^\infty(\Gamma \cup W)} \|\operatorname{tr}(\nabla N)\|_{L^\infty(\Omega^-)} \|\varpi\|_{L^2(\Omega^-)}^2 + C/\delta \|\theta\|_{L^\infty(\Gamma \cup W)} \|\varpi\|_{L^2(\Omega^-)}^2 \\
&\leq CK^2 \|\varpi\|_{L^2(\Omega^-)}^2 \leq CK^2 E_0(0).
\end{aligned}$$

We also have, with the help of (3.1), that

$$\|\varpi\|_{L^2(\Gamma \cup W)}^2 \leq CK \|\varpi\|_{L^2(\Omega^-)}^2 \leq CK E_0(0).$$

From Gauss' formula, Hölder's inequality and Cauchy's inequality, it follows

$$\begin{aligned}
\|\nabla^r \varpi\|_{L^2(\Gamma \cup W)}^2 &= \int_{\Gamma \cup W} N_a N^a |\nabla^r \varpi|^2 d\mu_\gamma = - \int_{\Omega^-} \nabla_a (N^a |\nabla^r \varpi|^2) d\mu_g \\
&\leq CK \|\nabla^r \varpi\|_{L^2(\Omega^-)}^2 + C \|\nabla^r \varpi\|_{L^2(\Omega^-)} \|\nabla^{r+1} \varpi\|_{L^2(\Omega^-)}. \quad (3.2)
\end{aligned}$$

Since $\operatorname{div} \varpi = 0$ and $\nabla^\perp \cdot \varpi = 0$, we have, from (A.6) and (A.7), that

$$\begin{aligned}
|\nabla^{r+1} \varpi|^2 &\leq C_1 g^{bc} \gamma^{af} \gamma^{AF} \nabla_A^r \nabla_a \varpi_b \nabla_F^r \nabla_f \varpi_c, \\
\int_{\Omega^-} |\nabla^{r+1} \varpi|^2 d\mu_g &\leq C_2 \int_{\Omega^-} (g^{bc} N^a N^f \gamma^{AF} \nabla_A^r \nabla_b \varpi_a \nabla_F^r \nabla_c \varpi_f + K^2 |\nabla^r \varpi|^2) d\mu_g.
\end{aligned}$$

Noticing that $\nabla^\perp \cdot \varpi = 0$, it follows from the Gauss formula that

$$\begin{aligned}
&(C_1^{-1} + C_2^{-1}) \|\nabla^{r+1} \varpi\|_{L^2(\Omega^-)}^2 \\
&\leq \int_{\Omega^-} g^{bc} \gamma^{af} \gamma^{AF} \nabla_A^r \nabla_a \varpi_b \nabla_F^r \nabla_c \varpi_f d\mu_g \\
&\quad + \int_{\Omega^-} (g^{bc} N^a N^f \gamma^{AF} \nabla_A^r \nabla_b \varpi_a \nabla_F^r \nabla_c \varpi_f + K^2 |\nabla^r \varpi|^2) d\mu_g \\
&= \int_{\Omega^-} g^{bc} \nabla_c (g^{af} \gamma^{AF} \nabla_A^r \nabla_a \varpi_b \nabla_F^r \varpi_f) d\mu_g \quad (3.3) \\
&\quad - \int_{\Omega^-} g^{bc} g^{af} \nabla_c (\gamma^{AF}) \nabla_A^r \nabla_a \varpi_b \nabla_F^r \varpi_f d\mu_g \quad (3.4) \\
&\quad + \int_{\Omega^-} K^2 |\nabla^r \varpi|^2 d\mu_g. \quad (3.5)
\end{aligned}$$

By the Hölder inequalities and the Cauchy inequality, we get

$$|(3.4)| \leq CK \|\nabla^r \varpi\|_{L^2(\Omega^-)} \|\nabla^{r+1} \varpi\|_{L^2(\Omega^-)}.$$

We write

$$(3.3) = - \int_{\Gamma \cup W} N^b g^{af} \gamma^{AF} \nabla_A^r \nabla_a \varpi_b \nabla_F^r \varpi_f d\mu_\gamma$$

$$= - \int_{\Gamma \cup W} \gamma^{af} \nabla_a (N^b \gamma^{AF} \nabla_A^r \varpi_b \nabla_F^r \varpi_f - N_f N^c N^b \gamma^{AF} \nabla_A^r \varpi_b \nabla_F^r \varpi_c) d\mu_\gamma \quad (3.6)$$

$$+ \int_{\Gamma \cup W} \gamma^{af} \nabla_a (N^b \gamma^{AF}) \nabla_A^r \varpi_b \nabla_F^r \varpi_f d\mu_\gamma \quad (3.7)$$

$$+ \int_{\Gamma \cup W} \gamma^{af} N^b \gamma^{AF} \nabla_A^r \varpi_b \nabla_F^r \nabla_a \varpi_f d\mu_\gamma \quad (3.8)$$

$$+ \int_{\Gamma \cup W} N^a N^f N^b \gamma^{AF} \nabla_A^r \varpi_b \nabla_F^r \nabla_a \varpi_f d\mu_\gamma \quad (3.9)$$

$$- \int_{\Gamma \cup W} \gamma^{af} \nabla_a N_f N^c N^b \gamma^{AF} \nabla_A^r \varpi_b \nabla_F^r \varpi_c d\mu_\gamma. \quad (3.10)$$

In view of the Gauss formula, (3.6) vanishes. Due to $\operatorname{div} \varpi = 0$, (3.8) + (3.9) = 0. From the Hölder inequality, we have

$$|(3.7) + (3.10)| \leq CK \|\nabla^r \varpi\|_{L^2(\Gamma \cup W)}^2.$$

Thus, from (3.2) and the Cauchy inequality, we get

$$\begin{aligned} \|\nabla^{r+1} \varpi\|_{L^2(\Omega^-)}^2 &\leq CK \|\nabla^r \varpi\|_{L^2(\Gamma \cup W)}^2 + CK \|\nabla^r \varpi\|_{L^2(\Omega^-)} \|\nabla^{r+1} \varpi\|_{L^2(\Omega^-)} \\ &\quad + CK^2 \|\nabla^r \varpi\|_{L^2(\Omega^-)}^2 \\ &\leq CK^2 \|\nabla^r \varpi\|_{L^2(\Omega^-)}^2 + CK \|\nabla^r \varpi\|_{L^2(\Omega^-)} \|\nabla^{r+1} \varpi\|_{L^2(\Omega^-)} \\ &\leq CK^2 \|\nabla^r \varpi\|_{L^2(\Omega^-)}^2 + \frac{1}{2} \|\nabla^{r+1} \varpi\|_{L^2(\Omega^-)}^2, \end{aligned}$$

namely,

$$\|\nabla^{r+1} \varpi\|_{L^2(\Omega^-)}^2 \leq CK^2 \|\nabla^r \varpi\|_{L^2(\Omega^-)}^2, \quad (3.11)$$

and then

$$\|\nabla^r \varpi\|_{L^2(\Gamma \cup W)}^2 \leq CK \|\nabla^r \varpi\|_{L^2(\Omega^-)}^2. \quad (3.12)$$

Therefore, by the induction argument, we have obtained the desired results. $\square$

Lemma 3.2. *Let α be a $(0, r)$ tensor defined in $\overline{\Omega^-} \subset \mathbb{R}^2$, and assume that $|\theta|_{L^\infty(\Gamma \cup W)} + 1/\iota_0 \leq K$ and $\operatorname{Vol} \Omega^- \leq V$. Then there is $C = C(K, V, r, p)$ such that*

$$\|\alpha\|_{L^{p/(2-p)}(\Gamma \cup W)} \leq C \|\nabla \alpha\|_{L^p(\Omega^-)} + C \|\alpha\|_{L^p(\Omega^-)}, \quad 1 \leq p < 2. \quad (3.13)$$

Proof. By Hölder's inequalities, Cauchy's inequality, we get

$$\begin{aligned} \|\alpha\|_{L^{p/(2-p)}(\Gamma \cup W)}^{p/(2-p)} &= \int_{\Gamma} |\alpha|^{p/(2-p)} d\mu_\gamma + \int_W |\alpha|^{p/(2-p)} d\mu_\gamma \\ &= - \int_{\Omega^-} \nabla_a (N^a |\alpha|^{p/(2-p)}) d\mu_g \\ &\leq CK \|\alpha\|_{L^{p/(2-p)}(\Omega^-)}^{p/(2-p)} + C \frac{p}{2-p} \|\alpha\|_{L^{p/(2-p)}(\Omega^-)}^{2(p-1)/(2-p)} \|\nabla \alpha\|_{L^{p/(2-p)}(\Omega^-)} \\ &\leq C \left(K + \frac{p}{2-p}\right) \|\alpha\|_{L^{p/(2-p)}(\Omega^-)}^{p/(2-p)} + C \frac{p}{2-p} \|\nabla \alpha\|_{L^{p/(2-p)}(\Omega^-)}^{p/(2-p)}. \end{aligned}$$

Thus, we have

$$\begin{aligned} \|\alpha\|_{L^{p/(2-p)}(\Gamma \cup W)} &\leq C(K, r, p) \|\alpha\|_{L^{p/(2-p)}(\Omega^-)} + C(r, p) \|\nabla \alpha\|_{L^{p/(2-p)}(\Omega^-)} \\ &\leq (\text{Vol } \Omega^-)^{(1-p)/p} (C(K, r, p) \|\alpha\|_{L^p(\Omega^-)} + C(r, p) \|\nabla \alpha\|_{L^p(\Omega^-)}) \\ &\leq C(K, V, r, p) (\|\nabla \alpha\|_{L^p(\Omega^-)} + \|\alpha\|_{L^p(\Omega^-)}). \end{aligned}$$

□

Proposition 3.3. *Suppose that for $\iota_1 \geq 1/K_1$,*

$$|N(\bar{x}_1) - N(\bar{x}_2)| \leq \varepsilon_1, \quad \text{whenever } |\bar{x}_1 - \bar{x}_2| \leq \iota_1, \quad \bar{x}_1, \bar{x}_2 \in \Gamma \cup W,$$

and

$$C_0^{-1} \gamma_{ab}^0(y) Z^a Z^b \leq \gamma_{ab}(t, y) Z^a Z^b \leq C_0 \gamma_{ab}^0(y) Z^a Z^b, \quad \text{if } Z \in T(\Omega^-),$$

where $\gamma_{ab}^0(y) = \gamma_{ab}(0, y)$. Then, it holds for any integer $r \geq 0$

$$\|\nabla^r \varpi\|_{L^\infty(\Gamma \cup W)} \leq C(r, K, K_1, \text{Vol } \Omega^-) E_0^{1/2}(0),$$

and

$$\|\Pi \nabla^r |\varpi|^2\|_{L^2(\Gamma \cup W)} \leq C(r, K, K_1, \text{Vol } \Omega^-) E_0(0).$$

Proof. From (A.10) and (3.13), it follows

$$\begin{aligned} \|\nabla^s \varpi\|_{L^\infty(\Gamma \cup W)} &\leq C \|\nabla^{s+1} \varpi\|_{L^4(\Gamma \cup W)} + C(K_1) \|\nabla^s \varpi\|_{L^4(\Gamma \cup W)} \\ &\leq C(K, K_1, \text{Vol } \Omega^-) \sum_{\ell=0}^2 \|\nabla^{s+\ell} \varpi\|_{L^2(\Omega^-)} \\ &\leq C(K, K_1, \text{Vol } \Omega^-) \sum_{\ell=0}^2 K^{s+\ell} E_0^{1/2}(0) \\ &\leq C(s, K, K_1, \text{Vol } \Omega^-) E_0^{1/2}(0). \end{aligned}$$

By the Hölder inequality, Proposition 3.1, (A.10), and the Cauchy inequality, we have

$$\begin{aligned} \|\Pi \nabla^r |\varpi|^2\|_{L^2(\Gamma \cup W)} &\leq C \sum_{m=0}^{[r/2]} \|\nabla^m \varpi\|_{L^\infty(\Gamma \cup W)} \|\nabla^{r-m} \varpi\|_{L^2(\Gamma \cup W)} \\ &\leq C \sum_{m=0}^{[r/2]} (\|\nabla^{m+1} \varpi\|_{L^4(\Gamma \cup W)} + C(K_1) \|\nabla^m \varpi\|_{L^4(\Gamma \cup W)}) K^{r-m+1/2} E_0^{1/2}(0) \\ &\leq C(K_1, \text{Vol } \Omega^-) E_0^{1/2}(0) \sum_{m=0}^{[r/2]} \sum_{\ell=0}^2 \|\nabla^{m+\ell} \varpi\|_{L^2(\Omega^-)} K^{r-m+1/2} \\ &\leq C(K_1, \text{Vol } \Omega^-) \sum_{m=0}^{[r/2]} \sum_{\ell=0}^2 K^{m+\ell} K^{r-m+1/2} E_0(0) \\ &\leq C(r, K, K_1, \text{Vol } \Omega^-) E_0(0). \end{aligned}$$

□

4. THE ESTIMATES OF THE ELECTRIC FIELD IN VACUUM

Although the electric field can be regarded as a secondary variable due to

$$\nabla^\perp \hat{E} = -\hat{\mathbf{H}}_t,$$

we have to use the estimates of the electric field in vacuum in order to get the energy estimates. In fact, we can prove the following estimates.

Proposition 4.1. *If $|\theta| + 1/\iota_0 \leq K$ on $\Gamma \cup W$ and $\nabla^\perp \Xi_0 \in L^2(\Omega^-)$, then it holds for any integer $r \geq 0$*

$$\begin{aligned} & \|\nabla^r(\nabla^\perp \Xi)\|_{L^2(\Omega^-)}^2 + \|\nabla^r(\nabla^\perp \Xi)\|_{L^2(\Gamma \cup W)}^2 \\ & \leq C(r, K) \left[\|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)}^2 + E_0(0) \sup_{\tau \in [0, t]} \sum_{\ell=0}^2 \|\nabla^\ell u(\tau)\|_{L^2(\Gamma)}^2 \right] \\ & \quad \cdot \exp \left(C(K) \int_0^t \|u_N(\tau)\|_{L^\infty(\Gamma)} d\tau \right), \end{aligned}$$

where Ξ_0 is the initial datum of Ξ at $t = 0$.

Proof. From (2.13c) and (A.4), we have on boundaries $\Gamma \cup W$

$$N^a \nabla_a^\perp \Xi = N^d u^a \nabla_a \varpi_d - \nabla_a u_d N^d \varpi^a.$$

It is clear to see that Ξ is a solution of the following mixed problem

$$\begin{cases} \Delta \Xi = 0, & \text{in } \Omega^-, \end{cases} \quad (4.1a)$$

$$\begin{cases} \Xi = f_1, & \nabla_{N^\perp} \Xi = f_2, & \text{on } \Gamma, \end{cases} \quad (4.1b)$$

$$\begin{cases} \Xi = 0, & \nabla_{N^\perp} \Xi = 0, & \text{on } W, \end{cases} \quad (4.1c)$$

where $f_1 = u_N(\varpi_1 N_2 - \varpi_2 N_1)$ and $f_2 = \nabla_a u_d N^d \varpi^a - N^d u^a \nabla_a \varpi_d$.

By Hölder's inequality, we get

$$\begin{aligned} & \|\overline{\nabla} \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 = \int_{\Omega^-} g^{ab} \gamma^{cd} \nabla_c \nabla_a^\perp \Xi \nabla_d \nabla_b^\perp \Xi d\mu_g \\ & = \int_{\Omega^-} g^{ab} \nabla_a^\perp (\gamma^{cd} \nabla_c \Xi \nabla_d \nabla_b \Xi) d\mu_g - \int_{\Omega^-} g^{ab} \nabla_a^\perp \gamma^{cd} \nabla_c \Xi \nabla_d \nabla_b \Xi d\mu_g \\ & = - \int_{\Gamma \cup W} (N^\perp)^b \gamma^{cd} \nabla_c \Xi \nabla_d \nabla_b \Xi d\mu_\gamma + \int_{\Omega^-} g^{ab} \nabla_a \gamma^{cd} \nabla_c \Xi \nabla_d \nabla_b^\perp \Xi d\mu_g \\ & = - \int_\Gamma \gamma^{cd} \nabla_c \Xi \nabla_d (\nabla_{N^\perp} \Xi) d\mu_\gamma - \int_\Gamma \gamma^{cd} \nabla_c \Xi \nabla_d N^b \nabla_b^\perp \Xi d\mu_\gamma \\ & \quad + \int_{\Omega^-} g^{ab} \nabla_a \gamma^{cd} \nabla_c \Xi \nabla_d \nabla_b^\perp \Xi d\mu_g \\ & \leq C \|\overline{\nabla} f_1\|_{L^2(\Gamma)} \|\overline{\nabla} f_2\|_{L^2(\Gamma)} + C \|\theta\|_{L^\infty(\Gamma)} \|\overline{\nabla} f_1\|_{L^2(\Gamma)} \|\nabla^\perp \Xi\|_{L^2(\Gamma)} \\ & \quad + CK \|\nabla^\perp \Xi\|_{L^2(\Omega^-)} \|\nabla \nabla^\perp \Xi\|_{L^2(\Omega^-)}. \end{aligned}$$

Similar to (3.1), we have for any $\delta > 0$

$$\|\nabla^\perp \Xi\|_{L^2(\Gamma)}^2 \leq C(K + 1/\delta) \|\nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 + \delta \|\nabla \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2. \quad (4.2)$$

Thus, in view of (A.6) and Cauchy's inequality, it follows that

$$\begin{aligned} & \|\nabla \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 \leq C \|\overline{\nabla} \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 \\ & \leq C \|\overline{\nabla} f_1\|_{L^2(\Gamma)} \|\overline{\nabla} f_2\|_{L^2(\Gamma)} + C \|\overline{\nabla} f_1\|_{L^2(\Gamma)}^2 \\ & \quad + C(K + 1)(K + 1/\delta) \|\nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 + C(K + 1)\delta \|\nabla \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 \end{aligned}$$

$$+ CK^2 \|\nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 + \frac{1}{4} \|\nabla \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2.$$

Taken δ so small that $C(K+1)\delta < 1/4$, it yields

$$\|\nabla \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 \leq C \|\bar{\nabla} f_1\|_{L^2(\Gamma)} \|\bar{\nabla} f_2\|_{L^2(\Gamma)} + C \|\bar{\nabla} f\|_{L^2(\Gamma)}^2 + C(K) \|\nabla^\perp \Xi\|_{L^2(\Omega^-)}^2. \quad (4.3)$$

Similar to the derivation of (2.13c), we can get, due to $\hat{E}_t = 0$, that

$$D_t \nabla_a^\perp \Xi = u^b \nabla_b \nabla_a^\perp \Xi + \nabla_a u^b \nabla_b^\perp \Xi. \quad (4.4)$$

It follows that

$$\frac{d}{dt} \int_{\Omega^-} |\nabla^\perp \Xi|^2 d\mu_g = - \int_{\Gamma} u_N |\nabla^\perp \Xi|^2 d\mu_\gamma,$$

which yields

$$\|\nabla^\perp \Xi(t)\|_{L^2(\Omega^-)}^2 \leq \|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)}^2 + \int_0^t \|u_N(\tau)\|_{L^\infty(\Gamma)} \|\nabla^\perp \Xi(\tau)\|_{L^2(\Gamma)}^2 d\tau. \quad (4.5)$$

From (4.3), (4.5) and (4.2), we obtain

$$\begin{aligned} & \|\nabla^\perp \Xi(t)\|_{L^2(\Omega^-)}^2 + \|\nabla \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 \\ & \leq C \|\bar{\nabla} f_1\|_{L^2(\Gamma)}^2 + C \|\bar{\nabla} f_2\|_{L^2(\Gamma)}^2 + C(K) \|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)}^2 \\ & \quad + C(K) \int_0^t \|u_N(\tau)\|_{L^\infty(\Gamma)} \|\nabla^\perp \Xi(\tau)\|_{L^2(\Gamma)}^2 d\tau \\ & \leq C \|\bar{\nabla} f_1\|_{L^2(\Gamma)}^2 + C \|\bar{\nabla} f_2\|_{L^2(\Gamma)}^2 + C(K) \|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)}^2 \\ & \quad + C(K) \int_0^t \|u_N(\tau)\|_{L^\infty(\Gamma)} (\|\nabla^\perp \Xi(\tau)\|_{L^2(\Omega^-)}^2 + \|\nabla \nabla^\perp \Xi(\tau)\|_{L^2(\Omega^-)}^2) d\tau, \end{aligned}$$

which implies, by Grönwall's inequality, that

$$\begin{aligned} & \|\nabla^\perp \Xi(t)\|_{L^2(\Omega^-)}^2 + \|\nabla \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 \\ & \leq \left[C \sup_{\tau \in [0, t]} [\|\bar{\nabla} f_1(\tau)\|_{L^2(\Gamma)}^2 + \|\bar{\nabla} f_2(\tau)\|_{L^2(\Gamma)}^2] + C(K) \|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)}^2 \right] \\ & \quad \cdot \exp \left(C(K) \int_0^t \|u_N(\tau)\|_{L^\infty(\Gamma)} d\tau \right). \end{aligned}$$

By the definition of f_1 and f_2 and Proposition 3.1, we have

$$\begin{aligned} \|\bar{\nabla} f_1\|_{L^2(\Gamma)} &= \|\bar{\nabla}[u_N(\varpi_1 N_2 - \varpi_2 N_1)]\|_{L^2(\Gamma)} \\ &\leq C \|\theta\|_{L^\infty(\Gamma)} \|u\|_{L^2(\Gamma)} \|\varpi\|_{L^2(\Gamma)} + \|\nabla u\|_{L^2(\Gamma)} \|\varpi\|_{L^2(\Gamma)} + \|u\|_{L^2(\Gamma)} \|\nabla \varpi\|_{L^2(\Gamma)} \\ &\leq C(K) [\|u\|_{L^2(\Gamma)} + \|\nabla u\|_{L^2(\Gamma)}] E_0^{1/2}(0), \\ \|\bar{\nabla} f_2\|_{L^2(\Gamma)} &= \|\bar{\nabla}(N^d u^a \nabla_a \varpi_d - \nabla_a u_d N^d \varpi^a)\|_{L^2(\Gamma)} \\ &\leq \|\theta\|_{L^\infty(\Gamma)} \|u\|_{L^2(\Gamma)} \|\nabla \varpi\|_{L^2(\Gamma)} + \|\nabla u\|_{L^2(\Gamma)} \|\nabla \varpi\|_{L^2(\Gamma)} \\ &\quad + \|u\|_{L^2(\Gamma)} \|\nabla^2 \varpi\|_{L^2(\Gamma)} + \|\nabla^2 u\|_{L^2(\Gamma)} \|\varpi\|_{L^2(\Gamma)} \\ &\quad + \|\theta\|_{L^\infty(\Gamma)} \|\nabla u\|_{L^2(\Gamma)} \|\varpi\|_{L^2(\Gamma)} \\ &\leq C(K) [\|u\|_{L^2(\Gamma)} + \|\nabla u\|_{L^2(\Gamma)} + \|\nabla^2 u\|_{L^2(\Gamma)}] E_0^{1/2}(0) \\ &\leq C(K) E_0^{1/2}(0) \sum_{\ell=0}^2 \|\nabla^\ell u\|_{L^2(\Gamma)}. \end{aligned}$$

Therefore, we obtain, combining with (4.2), that

$$\begin{aligned} & \|\nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 + \|\nabla \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 + \|\nabla^\perp \Xi\|_{L^2(\Gamma \cup W)}^2 \\ & \leq C(K) \left[\|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)}^2 + E_0(0) \sup_{\tau \in [0, t]} \sum_{\ell=0}^2 \|\nabla^\ell u(\tau)\|_{L^2(\Gamma)}^2 \right] \\ & \quad \cdot \exp \left(C(K) \int_0^t \|u_N(\tau)\|_{L^\infty(\Gamma)} d\tau \right). \end{aligned}$$

Since $\nabla_a(\nabla^\perp \Xi)^a = 0$ and $\nabla^\perp \cdot \nabla^\perp \Xi = 0$, it is similar to ϖ . One can verify that the lines between (3.2) and (3.12) also hold if ϖ is replaced by $\nabla^\perp \Xi$ everywhere in those lines. Thus, we can obtain for any $r \geq 1$

$$\|\nabla^{r+1} \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 \leq CK^2 \|\nabla^r \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2, \quad (4.6)$$

and

$$\|\nabla^r \nabla^\perp \Xi\|_{L^2(\Gamma \cup W)}^2 \leq CK \|\nabla^r \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2. \quad (4.7)$$

Hence, we get for any $r \geq 0$

$$\begin{aligned} & \|\nabla^r \nabla^\perp \Xi\|_{L^2(\Omega^-)}^2 + \|\nabla^r \nabla^\perp \Xi\|_{L^2(\Gamma \cup W)}^2 \\ & \leq C(K, \text{Vol } \Omega^+) \left[\|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)}^2 + E_0(0) \sup_{\tau \in [0, t]} \sum_{\ell=0}^2 \|\nabla^\ell u(\tau)\|_{L^2(\Omega^+)}^2 \right] \\ & \quad \cdot \exp \left(C(K) \int_0^t \|u_N(\tau)\|_{L^\infty(\Gamma)} d\tau \right). \end{aligned}$$

Thus, we obtain the desired results. $\square$

5. THE FIRST ORDER ENERGY ESTIMATES

We have

$$D_t \nabla_a u_b + \nabla_a \nabla_b q^+ = \nabla_b u_c \nabla_a u^c + \mu \beta^c \nabla_c \nabla_a \beta_b + \mu \nabla_a \beta^c \nabla_c \beta_b, \quad (5.1)$$

$$D_t \nabla_a \beta_b = \nabla_b u_c \nabla_a \beta^c + \beta^c \nabla_c \nabla_a u_b + \nabla_a \beta^c \nabla_c u_b. \quad (5.2)$$

The first order energy is given by

$$\begin{aligned} E_1(t) &= \int_{\Omega^+} g^{bd} \gamma^{af} \nabla_a u_b \nabla_f u_d d\mu_g + \mu \int_{\Omega^+} g^{bd} \gamma^{af} \nabla_a \beta_b \nabla_f \beta_d d\mu_g \\ & \quad + \int_{\Omega^+} |\nabla^\perp \cdot u|^2 d\mu_g + \mu \int_{\Omega^+} |\nabla^\perp \cdot \beta|^2 d\mu_g. \end{aligned}$$

Then, we have the following estimates.

Proposition 5.1. *If for any smooth solutions of (2.13), it holds $\|\nabla p(t)\|_{L^2(\Omega^+)} \leq M$ in $[0, T]$ and*

$$|\nabla u| \leq M, \quad \text{in } [0, T] \times \Omega^+, \quad (5.3)$$

$$|\theta| + 1/\iota_0 \leq K, \quad \text{on } [0, T] \times \Gamma, \quad (5.4)$$

then, we have for $t \in [0, T]$

$$E_1(t) \leq e^{(C(K)M+1)t} E_1(0) + C(K, K_1, \text{Vol } \Omega) E_0(0) (e^{(C(K)M+1)t} - 1). \quad (5.5)$$

Proof. Since $D_t d\mu_g = \text{tr } h d\mu_g = 0$ due to $\text{tr } h = \nabla_a u^a = 0$, we have, by noticing that $\nabla_a \beta^a = 0$ in Ω^+ , that

$$\begin{aligned}
\frac{d}{dt} E_1(t) &= \int_{\Omega^+} D_t g^{bd} \gamma^{af} \nabla_a u_b \nabla_f u_d d\mu_g + \int_{\Omega^+} g^{bd} D_t \gamma^{af} \nabla_a u_b \nabla_f u_d d\mu_g \\
&\quad + 2 \int_{\Omega^+} g^{bd} \gamma^{af} D_t \nabla_a u_b \nabla_f u_d d\mu_g + \mu \int_{\Omega^+} D_t g^{bd} \gamma^{af} \nabla_a \beta_b \nabla_f \beta_d d\mu_g \\
&\quad + \mu \int_{\Omega^+} g^{bd} D_t \gamma^{af} \nabla_a \beta_b \nabla_f \beta_d d\mu_g + 2\mu \int_{\Omega^+} g^{bd} \gamma^{af} D_t \nabla_a \beta_b \nabla_f \beta_d d\mu_g \\
&\quad + 2\mu \int_{\Omega^-} g^{bd} \gamma^{af} D_t \nabla_a \varpi_b \nabla_f \varpi_d d\mu_g \\
&\quad + 2 \int_{\Omega^+} (\nabla_1 u_2 - \nabla_2 u_1) D_t (\nabla_1 u_2 - \nabla_2 u_1) d\mu_g \\
&\quad + 2\mu \int_{\Omega^+} (\nabla_1 \beta_2 - \nabla_2 \beta_1) D_t (\nabla_1 \beta_2 - \nabla_2 \beta_1) d\mu_g \\
&= -2 \int_{\Omega^+} h^{bd} \gamma^{af} \nabla_a u_b \nabla_f u_d d\mu_g - 2 \int_{\Omega^+} \gamma^{ac} h_{ce} \gamma^{ef} \nabla_a u_b \nabla_f u^b d\mu_g \\
&\quad - 2 \int_{\Gamma} N_b \gamma^{af} \nabla_a q^+ \nabla_f u^b d\mu_\gamma + 2 \int_{\Omega^+} \nabla_b \gamma^{af} \nabla_a q^+ \nabla_f u^b d\mu_g \\
&\quad + 2 \int_{\Omega^+} \gamma^{af} \nabla_b u_c \nabla_a u^c \nabla_f u^b d\mu_g + 2\mu \int_{\Gamma} N_c \beta^c \gamma^{af} \nabla_a \beta_b \nabla_f u^b d\mu_\gamma \\
&\quad - 2\mu \int_{\Omega^+} \nabla_c \gamma^{af} \beta^c \nabla_a \beta_b \nabla_f u^b d\mu_g + 2\mu \int_{\Omega^+} \gamma^{af} \nabla_a \beta^c \nabla_c \beta_b \nabla_f u^b d\mu_g \\
&\quad - 2\mu \int_{\Omega^+} h^{bd} \gamma^{af} \nabla_a \beta_b \nabla_f \beta_d d\mu_g - 2\mu \int_{\Omega^+} \gamma^{ac} h_{ce} \gamma^{ef} \nabla_a \beta_b \nabla_f \beta^b d\mu_g \\
&\quad + 2\mu \int_{\Omega^+} \gamma^{af} \nabla_b u_c \nabla_a \beta^c \nabla_f \beta^b d\mu_g + 2\mu \int_{\Omega^+} \gamma^{af} \nabla_a \beta^c \nabla_c u_b \nabla_f \beta^b d\mu_g \\
&\quad + 2\mu \int_{\Gamma} N_c \beta^c (\nabla_1 u_2 - \nabla_2 u_1) (\nabla_1 \beta_2 - \nabla_2 \beta_1) d\mu_\gamma \\
&\quad + 4\mu \int_{\Omega^+} (\nabla_1 \beta_2 - \nabla_2 \beta_1) \nabla_1 \beta_1 (\nabla_1 u_2 + \nabla_2 u_1) d\mu_g \\
&\quad + 4\mu \int_{\Omega^+} (\nabla_1 \beta_2 - \nabla_2 \beta_1) \nabla_2 u_2 (\nabla_1 \beta_2 + \nabla_2 \beta_1) d\mu_g.
\end{aligned}$$

From (2.13f), the last two boundary integrals vanish, and the first boundary integral above is equal to

$$\begin{aligned}
&-2 \int_{\Gamma} N_b \gamma^{af} \nabla_a q^- \nabla_f u^b d\mu_\gamma \\
&= -2 \int_{\Gamma} \gamma^{af} \nabla_f (N_b \nabla_a q^- u^b - N_a N^c N_b \nabla_c q^- u^b) d\mu_\gamma
\end{aligned} \tag{5.6}$$

$$+ 2 \int_{\Gamma} \text{tr } \theta N^c N_b \nabla_c q^- u^b d\mu_\gamma + 2 \int_{\Gamma} \gamma^{af} \nabla_f N_b \nabla_a q^- u^b d\mu_\gamma \tag{5.7}$$

$$+ 2 \int_{\Gamma} \gamma^{af} N_b \nabla_f \nabla_a q^- u^b d\mu_\gamma \tag{5.8}$$

$$\begin{aligned}
&\leq C\mu \|\theta\|_{L^\infty(\Gamma)} \|\varpi\|_{L^2(\Gamma)} \|\nabla \varpi\|_{L^2(\Gamma)} \|u\|_{L^\infty(\Gamma)} + C\mu \|\Pi \nabla^2 |\varpi|^2\|_{L^2(\Gamma)} \|u\|_{L^2(\Gamma)} \\
&\leq C(K, K_1, \text{Vol } \Omega) (E_0(0) + E_0^{1/2}(0) E_1^{1/2}(t)),
\end{aligned}$$

where (5.6) vanishes due to Gauss' formula, and we have used Propositions 3.1 and 3.3, and $\|u\|_{L^2(\Gamma)} \leq C(K, \text{Vol } \Omega^+)(\|u\|_{L^2(\Omega^+)} + \|\nabla u\|_{L^2(\Omega^+)})$ in view of (3.13).

Thus, by Hölder's inequalities and $|\nabla u| + |\nabla p| \leq M$ in Ω^+ , we get

$$\frac{d}{dt}E_1(t) \leq C(K, K_1, \text{Vol } \Omega)E_0(0) + (C(K)M + 1)E_1(t),$$

which yields (5.5) from the Grönwall inequality. $\square$

6. THE SECOND ORDER ENERGY ESTIMATES

By (2.7), we have

$$\begin{aligned} D_t \nabla_{a_2} \nabla_{a_1} u_b &= -\nabla_{a_2} \nabla_{a_1} \nabla_b q^+ + \nabla_{a_2} \nabla_{a_1} u^c (\nabla_b u_c - \nabla_c u_b) + \mu \nabla_{a_2} \beta^c \nabla_c \nabla_{a_1} \beta_b \\ &\quad + \mu \beta^c \nabla_c \nabla_{a_2} \nabla_{a_1} \beta_b + \mu \nabla_{a_2} \nabla_{a_1} \beta^c \nabla_c \beta_b + \mu \nabla_{a_1} \beta^c \nabla_{a_2} \nabla_c \beta_b, \end{aligned} \quad (6.1)$$

$$\begin{aligned} D_t \nabla_{a_2} \nabla_{a_1} \beta_b &= -\nabla_{a_2} \nabla_{a_1} u^c \nabla_c \beta_b + h_{bc} \nabla_{a_2} \nabla_{a_1} \beta^c + \beta^c \nabla_c \nabla_{a_2} \nabla_{a_1} u_b \\ &\quad + \nabla_{a_1} \beta^c \nabla_{a_2} \nabla_c u_b + \nabla_{a_2} \beta^c \nabla_{a_1} \nabla_c u_b. \end{aligned} \quad (6.2)$$

Define the r -th order energy as

$$\begin{aligned} E_2(t) &= \int_{\Omega^+} g^{bd} \gamma^{a_1 f_1} \gamma^{a_2 f_2} \nabla_{a_2} \nabla_{a_1} u_b \nabla_{f_2} \nabla_{f_1} u_d d\mu_g \\ &\quad + \mu \int_{\Omega^+} g^{bd} \gamma^{a_1 f_1} \gamma^{a_2 f_2} \nabla_{a_2} \nabla_{a_1} \beta_b \nabla_{f_2} \nabla_{f_1} \beta_d d\mu_g \\ &\quad + \int_{\Omega^+} |\nabla \nabla^\perp \cdot u|^2 d\mu_g + \mu \int_{\Omega^+} |\nabla \nabla^\perp \cdot \beta|^2 d\mu_g \\ &\quad + \int_{\Gamma} \gamma^{a_1 f_1} \gamma^{a_2 f_2} \nabla_{a_2} \nabla_{a_1} P \nabla_{f_2} \nabla_{f_1} P \vartheta d\mu_\gamma, \end{aligned}$$

where $\vartheta = -1/\nabla_N P$.

Proposition 6.1. *For $t \in [0, T]$, assume that $|\theta| + 1/\iota_0 \leq K$ and $|\nabla_N D_t P| \leq L$ on Γ , $|\nabla u| + |\nabla \beta| \leq M$ in $\Omega^+ \cup \Gamma$, then we have for $t \in [0, T]$,*

$$E_2(t) \leq e^{Ct} E_2(0) + C(e^{Ct} - 1),$$

where the constants C 's depend on $K, K_1, M, L, 1/\varepsilon, \text{Vol } \Omega, E_0(0)$ and $E_1(0)$ and maybe differ from each other.

Proof. By (6.1) and $\text{div } u = 0$, we get

$$\begin{aligned} D_t \nabla_a (\nabla^\perp \cdot \beta) &= D_t \nabla_a (g^{de} \nabla_d^\perp \beta_e) = -D_t (g^{de} \nabla_a \nabla_d \beta_e^\perp) \\ &= 2h^{de} \nabla_a \nabla_d \beta_e^\perp - g^{de} D_t \nabla_a \nabla_d \beta_e^\perp \\ &= 2h^{de} \nabla_a \nabla_d \beta_e^\perp + g^{de} \nabla_a \nabla_d u^c \nabla_c \beta_e^\perp - \frac{1}{2} g^{de} (\nabla_e^\perp u_c + \nabla_c u_e^\perp) \nabla_a \nabla_d \beta^c \\ &\quad + \beta^c \nabla_c \nabla_a (\nabla^\perp \cdot u) - g^{de} \nabla_d \beta^c \nabla_a \nabla_c u_e^\perp + \nabla_a \beta^c \nabla_c (\nabla^\perp \cdot u), \end{aligned} \quad (6.3)$$

and similarly

$$\begin{aligned} D_t \nabla_a (\nabla^\perp \cdot u) &= 2h^{de} \nabla_a \nabla_d u_e^\perp - g^{de} \nabla_a \nabla_d u^c (\nabla_e^\perp u_c - \nabla_c u_e^\perp) + \mu \nabla_a \beta^c \nabla_c (\nabla^\perp \cdot \beta) \\ &\quad + \mu \beta^c \nabla_c \nabla_a (\nabla^\perp \cdot \beta) - \mu g^{de} \nabla_a \nabla_d \beta^c \nabla_c \beta_e^\perp - \mu g^{de} \nabla_d \beta^c \nabla_c \nabla_a \beta_e^\perp. \end{aligned} \quad (6.4)$$

Since $\text{tr } h = 0$, it follows that

$$\frac{d}{dt} \int_{\Omega^+} |\nabla \nabla^\perp \cdot u|^2 d\mu_g + \mu \frac{d}{dt} \int_{\Omega^+} |\nabla \nabla^\perp \cdot \beta|^2 d\mu_g$$

$$\begin{aligned}
&= \int_{\Omega^+} D_t[g^{ab}\nabla_a(\nabla^\perp \cdot u)\nabla_b(\nabla^\perp \cdot u)]d\mu_g + \mu \int_{\Omega^+} D_t[g^{ab}\nabla_a(\nabla^\perp \cdot \beta)\nabla_b(\nabla^\perp \cdot \beta)]d\mu_g \\
&= -2 \int_{\Omega^+} h^{ab}\nabla_a(\nabla^\perp \cdot u)\nabla_b(\nabla^\perp \cdot u)d\mu_g - 2\mu \int_{\Omega^+} h^{ab}\nabla_a(\nabla^\perp \cdot \beta)\nabla_b(\nabla^\perp \cdot \beta)d\mu_g \\
&\quad + 2 \int_{\Omega^+} g^{ab}D_t[\nabla_a(\nabla^\perp \cdot u)]\nabla_b(\nabla^\perp \cdot u)d\mu_g \\
&\quad + 2\mu \int_{\Omega^+} g^{ab}D_t[\nabla_a(\nabla^\perp \cdot \beta)]\nabla_b(\nabla^\perp \cdot \beta)d\mu_g \\
&= -2 \int_{\Omega^+} h^{ab}\nabla_a(\nabla^\perp \cdot u)\nabla_b(\nabla^\perp \cdot u)d\mu_g - 2\mu \int_{\Omega^+} h^{ab}\nabla_a(\nabla^\perp \cdot \beta)\nabla_b(\nabla^\perp \cdot \beta)d\mu_g \quad (6.5)
\end{aligned}$$

$$+ 4 \int_{\Omega^+} g^{ab}h^{de}\nabla_a\nabla_d u_e^\perp \nabla_b(\nabla^\perp \cdot u)d\mu_g \quad (6.6)$$

$$- 2 \int_{\Omega^+} g^{ab}g^{de}\nabla_a\nabla_d u^c(\nabla_e^\perp u_c - \nabla_c u_e^\perp)\nabla_b(\nabla^\perp \cdot u)d\mu_g \quad (6.7)$$

$$+ 2\mu \int_{\Omega^+} g^{ab}\nabla_a\beta^c\nabla_c(\nabla^\perp \cdot \beta)\nabla_b(\nabla^\perp \cdot u)d\mu_g \quad (6.8)$$

$$+ 2\mu \int_{\Gamma} N_c\beta^c g^{ab}\nabla_a(\nabla^\perp \cdot \beta)\nabla_b(\nabla^\perp \cdot u)d\mu_\gamma \quad (6.9)$$

$$- 2\mu \int_{\Omega^+} g^{ab}g^{de}\nabla_a\nabla_d\beta^c\nabla_c\beta_e^\perp \nabla_b(\nabla^\perp \cdot u)d\mu_g \quad (6.10)$$

$$- 2\mu \int_{\Omega^+} g^{ab}g^{de}\nabla_a\beta^c\nabla_c\nabla_d\beta_e^\perp \nabla_b(\nabla^\perp \cdot u)d\mu_g \quad (6.11)$$

$$+ 4\mu \int_{\Omega^+} g^{ab}h^{de}\nabla_a\nabla_d\beta_e^\perp \nabla_b(\nabla^\perp \cdot \beta)d\mu_g \quad (6.12)$$

$$+ 2\mu \int_{\Omega^+} g^{ab}g^{de}\nabla_a\nabla_d u^c\nabla_c\beta_e^\perp \nabla_b(\nabla^\perp \cdot \beta)d\mu_g \quad (6.13)$$

$$- \mu \int_{\Omega^+} g^{ab}g^{de}(\nabla_e^\perp u_c + \nabla_c u_e^\perp)\nabla_a\nabla_d\beta^c\nabla_b(\nabla^\perp \cdot \beta)d\mu_g \quad (6.14)$$

$$- 2\mu \int_{\Omega^+} g^{ab}g^{de}\nabla_a\beta^c\nabla_c\nabla_d u_e^\perp \nabla_b(\nabla^\perp \cdot \beta)d\mu_g \quad (6.15)$$

$$+ 2\mu \int_{\Omega^+} g^{ab}\nabla_a\beta^c\nabla_c(\nabla^\perp \cdot u)\nabla_b(\nabla^\perp \cdot \beta)d\mu_g. \quad (6.16)$$

Due to $\beta \cdot N = 0$ on Γ , (6.9) = 0. Thus, by the Hölder inequalities and the Gauss formula, we obtain

$$\begin{aligned}
&\frac{d}{dt} \int_{\Omega^+} |\nabla \nabla^\perp \cdot u|^2 d\mu_g + \mu \frac{d}{dt} \int_{\Omega^+} |\nabla \nabla^\perp \cdot \beta|^2 d\mu_g \\
&\leq C(K)(\|\nabla u\|_{L^\infty(\Gamma)} + \mu\|\nabla \beta\|_{L^\infty(\Gamma)})E_2(t) \leq C(K, M)E_2(t).
\end{aligned}$$

Therefore, by the Hölder inequalities, we have

$$\frac{d}{dt} E_2(t) \leq -2 \int_{\Omega^+} \gamma^{a_1 f_1} \gamma^{a_2 f_2} \nabla_{a_2} \nabla_{a_1} \nabla_b q^+ \nabla_{f_2} \nabla_{f_1} u^b d\mu_g \quad (6.17)$$

$$+ 2\mu \int_{\Gamma} N_c \beta^c \gamma^{a_1 f_1} \gamma^{a_2 f_2} \nabla_{a_2} \nabla_{a_1} \beta_b \nabla_{f_2} \nabla_{f_1} u^b d\mu_\gamma \quad (6.18)$$

$$+ 2 \int_{\Gamma} \gamma^{a_1 f_1} \gamma^{a_2 f_2} \nabla_{a_2} \nabla_{a_1} P D_t \nabla_{f_2} \nabla_{f_1} P \vartheta d\mu_\gamma \quad (6.19)$$

$$\begin{aligned}
& + \int_{\Gamma} \gamma^{a_1 f_1} \gamma^{a_2 f_2} \nabla_{a_2} \nabla_{a_1} P \nabla_{f_2} \nabla_{f_1} P \left(\frac{\vartheta_t}{\vartheta} - h_{NN} \right) \vartheta d\mu_{\gamma} \\
& + C(K, M) E_2(t),
\end{aligned} \tag{6.20}$$

where (6.18) = 0 because $\beta \cdot N = 0$ on the boundary Γ . From Gauss' formula, it follows that

$$(6.17) = -2 \int_{\Gamma} \gamma^{a_1 f_1} \gamma^{a_2 f_2} \nabla_{a_2} \nabla_{a_1} q^+ \nabla_{f_2} \nabla_{f_1} u^b N_b d\mu_{\gamma} \tag{6.21}$$

$$+ 2 \int_{\Omega^+} \nabla_b (\gamma^{a_1 f_1} \gamma^{a_2 f_2}) \nabla_{a_2} \nabla_{a_1} q^+ \nabla_{f_2} \nabla_{f_1} u^b d\mu_g. \tag{6.22}$$

By (2.11) and $\nabla_a P = N_a \nabla_N P$ (since $P = 0$ on Γ), we get

$$D_t \nabla^2 P = -\nabla^2 u \cdot \nabla P + \nabla^2 D_t P = (\nabla^2 u \cdot N) \vartheta^{-1} + \nabla^2 D_t P,$$

and then

$$(6.19) + (6.21) = 2 \int_{\Gamma} \gamma^{a_1 f_1} \gamma^{a_2 f_2} \nabla_{a_2} \nabla_{a_1} P \nabla_{f_2} \nabla_{f_1} D_t P \vartheta d\mu_{\gamma} \tag{6.23}$$

$$- 2 \int_{\Gamma} \gamma^{a_2 f_2} \gamma^{a_1 f_1} \nabla_{a_2} \nabla_{a_1} q^- \nabla_{f_2} \nabla_{f_1} u^b N_b d\mu_{\gamma}. \tag{6.24}$$

From Gauss' formula, Hölder's inequality, Proposition 3.1 and (A.13), it follows that

$$\begin{aligned}
(6.24) & = -2 \int_{\Gamma} \gamma^{a_2 f_2} \nabla_{f_2} [\gamma^{a_1 f_1} \nabla_{a_2} \nabla_{a_1} q^- \nabla_{f_1} u^b N_b - \gamma^{a_1 f_1} N_{a_2} N^c \nabla_c \nabla_{a_1} q^- \nabla_{f_1} u^b N_b] d\mu_{\gamma} \\
& + 2 \int_{\Gamma} \gamma^{a_2 f_2} \nabla_{f_2} [\gamma^{a_1 f_1} \nabla_{a_2} \nabla_{a_1} q^- N_b] \nabla_{f_1} u^b d\mu_{\gamma} \\
& - 2 \int_{\Gamma} \gamma^{a_2 f_2} \nabla_{f_2} N_{a_2} \gamma^{a_1 f_1} N^c \nabla_c \nabla_{a_1} q^- \nabla_{f_1} u^b N_b d\mu_{\gamma} \\
& \leq C(K) E_0(0) \|\nabla u\|_{L^2(\Gamma)} \\
& \leq C(K, \text{Vol } \Omega^+) E_0(0) (\|\nabla^2 u\|_{L^2(\Omega^+)} + \|\nabla u\|_{L^2(\Omega^+)}) \\
& \leq C(K, \text{Vol } \Omega^+) E_0(0) (E_2^{1/2}(t) + E_1^{1/2}(t)).
\end{aligned}$$

By Hölder's inequality, we get

$$(6.23) \leq C \|\vartheta\|_{L^\infty(\Gamma)}^{1/2} E_2^{1/2}(t) \|\Pi \nabla^2 D_t P\|_{L^2(\Gamma)},$$

$$(6.22) \leq C K \|\nabla^2 q^+\|_{L^2(\Omega^+)} E_2^{1/2}(t),$$

$$(6.20) \leq \frac{C}{\varepsilon} \|\nabla_N D_t P\|_{L^\infty(\Gamma)} E_2(t) \leq C(L/\varepsilon) E_2(t),$$

since from (A.4) and $\nabla_a P = N_a \nabla_N P$ on Γ

$$\frac{\vartheta_t}{\vartheta} = - \frac{D_t \nabla_N P}{\nabla_N P} = \frac{2h_d^a N^d \nabla_a P}{\nabla_N P} - h_{NN} - \frac{\nabla_N D_t P}{\nabla_N P} = h_{NN} - \frac{\nabla_N D_t P}{\nabla_N P}. \tag{6.25}$$

From (2.13), it follows that

$$\Delta q^+ = -\nabla_a u^b \nabla_b u^a + \mu \nabla_a \beta^b \nabla_b \beta^a. \tag{6.26}$$

By (A.8), (6.26), (A.2), Proposition 3.3, (A.11) and Hölder's inequality, we have

$$\begin{aligned}
& \|\nabla q^+\|_{L^2(\Gamma)} + \|\nabla^2 q^+\|_{L^2(\Omega^+)} \\
& \leq \delta \|\Pi \nabla^2 q^+\|_{L^2(\Gamma)} + C(1/\delta, K, \text{Vol } \Omega^+) \|\Delta q^+\|_{L^2(\Omega^+)} \\
& \leq \delta \|\Pi \nabla^2 P\|_{L^2(\Gamma)} + \delta \|\Pi \nabla^2 q^-\|_{L^2(\Gamma)}
\end{aligned}$$

$$\begin{aligned}
& + C(1/\delta, K, \text{Vol } \Omega^+, M)[\|\nabla u\|_{L^2(\Omega^+)} + \|\nabla \beta\|_{L^2(\Omega^+)}] \\
& \leq \delta \|\theta \nabla_N P\|_{L^2(\Gamma)} + \delta C(K, K_1, \text{Vol } \Omega^-) E_0(0) \\
& \quad + C(1/\delta, K, \text{Vol } \Omega^+, M)(K_1 \|\nabla u\|_{L^2(\Omega^+)} + \|\nabla^2 u\|_{L^2(\Omega^+)}) \\
& \quad + C(1/\delta, K, \text{Vol } \Omega^+, M)(K_1 \|\nabla \beta\|_{L^2(\Omega^+)} + \|\nabla^2 \beta\|_{L^2(\Omega^+)}) \\
& \leq \delta \|\theta\|_{L^\infty(\Gamma)} (\|\nabla q^+\|_{L^2(\Gamma)} + \|\nabla q^-\|_{L^2(\Gamma)}) + \delta C(K, K_1, \text{Vol } \Omega^-) E_0(0) \\
& \quad + C(1/\delta, K, K_1, \text{Vol } \Omega^+, M) E_1^{1/2}(t) + C(1/\delta, K, \text{Vol } \Omega^+, M) E_2^{1/2}(t) \\
& \leq \delta K \|\nabla q^+\|_{L^2(\Gamma)} + \delta C(K, K_1, \text{Vol } \Omega^-) E_0(0) \\
& \quad + C(1/\delta, K, K_1, \text{Vol } \Omega^+, M) E_1^{1/2}(t) + C(1/\delta, K, \text{Vol } \Omega^+, M) E_2^{1/2}(t),
\end{aligned}$$

which yields by taking $\delta K \leq 1/2$

$$\begin{aligned}
& \|\nabla q^+\|_{L^2(\Gamma)} + \|\nabla^2 q^+\|_{L^2(\Omega^+)} \\
& \leq C(K, K_1, \text{Vol } \Omega^-) E_0(0) + C(K, K_1, M, \text{Vol } \Omega^+) E_1^{1/2}(t) + C(K, M, \text{Vol } \Omega^+) E_2^{1/2}(t).
\end{aligned}$$

Thus, it implies, from Cauchy's inequality, that

$$(6.22) \leq C(K, K_1, \text{Vol } \Omega^-) E_0^2(0) + C(K, K_1, M, \text{Vol } \Omega^+) E_1(t) + C(K, M, \text{Vol } \Omega^+) E_2(t).$$

From (A.2) with the fact $D_t P = P = 0$ on Γ and Hölder's inequality, it follows that

$$\begin{aligned}
\|\Pi \nabla^2 D_t P\|_{L^2(\Gamma)} &= \|\theta \nabla_N D_t P\|_{L^2(\Gamma)} = \|\vartheta \Pi \nabla^2 P \nabla_N D_t P\|_{L^2(\Gamma)} \\
&\leq C \|\vartheta\|_{L^\infty(\Gamma)}^{1/2} \|\vartheta^{1/2} \Pi \nabla^2 P\|_{L^2(\Gamma)} \|\nabla_N D_t P\|_{L^\infty(\Gamma)} \\
&\leq C(1/\varepsilon) L E_2^{1/2}(t),
\end{aligned}$$

which yields

$$(6.23) \leq C(1/\varepsilon) L E_2(t).$$

Therefore,

$$\begin{aligned}
\frac{d}{dt} E_2(t) &\leq C(K, K_1, \text{Vol } \Omega^-) E_0^2(0) + C(K, K_1, M, \text{Vol } \Omega^+) E_1(t) \\
&\quad + C(1/\varepsilon, K, M, L, \text{Vol } \Omega^+) E_2(t),
\end{aligned}$$

which implies the desired estimates in view of the Grönwall inequality. $\square$

7. THE THIRD ORDER ENERGY ESTIMATES

We can get

$$\begin{aligned}
D_t \nabla^3 u_a + \nabla^3 \nabla_a q^+ &= (\nabla_a u_c - \nabla_c u_a) \nabla^3 u^c + \mu \beta^c \nabla_c \nabla^3 \beta_a + 3\mu \nabla \beta \cdot \nabla^3 \beta_a \\
&\quad + \mu \nabla^3 \beta^c \nabla_c \beta_a - 3\nabla^2 u^c \nabla \nabla_c u_a + 3\mu \nabla^2 \beta^c \nabla \nabla_c \beta_a,
\end{aligned} \tag{7.1}$$

and

$$\begin{aligned}
D_t \nabla^3 \beta_a &= (\nabla_a u_c + \nabla_c u_a) \nabla^3 \beta^c - \nabla^3 u^c \nabla_c \beta_a + \beta^c \nabla_c \nabla^3 u_a + 3\nabla \beta \cdot \nabla^3 u_a \\
&\quad - 3\nabla^2 u^c \nabla \nabla_c \beta_a + 3\nabla^2 \beta^c \nabla \nabla_c u_a.
\end{aligned} \tag{7.2}$$

Define the third order energy as

$$\begin{aligned}
E_3(t) &= \int_{\Omega^+} g^{bd} \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a u_b \nabla_F^2 \nabla_f u_d d\mu_g + \mu \int_{\Omega^+} g^{bd} \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a \beta_b \nabla_F^2 \nabla_f \beta_d d\mu_g \\
&\quad + \int_{\Omega^+} |\nabla^2(\nabla^\perp \cdot u)|^2 d\mu_g + \mu \int_{\Omega^+} |\nabla^2(\nabla^\perp \cdot \beta)|^2 d\mu_g
\end{aligned}$$

$$+ \int_{\Gamma} \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a P \nabla_F^2 \nabla_f P \vartheta d\mu_{\gamma},$$

where $\vartheta = -1/\nabla_N P$ as before.

Theorem 7.1. *For any smooth solution of MHD (2.13) satisfying*

$$|\nabla P| + |\nabla u| + |\nabla \beta| \leq M, \quad \text{in } [0, T] \times \Omega, \quad (7.3)$$

$$|\theta| + 1/\iota_0 \leq K, \quad \text{on } [0, T] \times \Gamma, \quad (7.4)$$

$$-\nabla_N P \geq \varepsilon > 0, \quad \text{on } [0, T] \times \Gamma, \quad (7.5)$$

$$|\nabla^2 P| + |\nabla_N D_t P| \leq L, \quad \text{on } [0, T] \times \Gamma, \quad (7.6)$$

we have, for $t \in [0, T]$,

$$E_3(t) \leq e^{C_1 t} E_3(0) + C_2 e^{C_3 t} (e^{C_4 t} - 1), \quad (7.7)$$

where the constant $C_i > 0$ depends on $K, K_1, M, M_1, L, 1/\varepsilon, \text{Vol } \Omega, E_0(0), E_1(0)$, and $E_2(0)$; both C_2 and C_3 also depend on $\|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)}$.

Proof. Since $\text{tr } h = 0$, we have

$$\frac{d}{dt} E_3(t) = \int_{\Omega^+} D_t (g^{bd} \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a u_b \nabla_F^2 \nabla_f u_d) d\mu_g \quad (7.8)$$

$$+ \mu \int_{\Omega^+} D_t (g^{bd} \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a \beta_b \nabla_F^2 \nabla_f \beta_d) d\mu_g \quad (7.9)$$

$$+ \int_{\Omega^+} D_t |\nabla^2 (\nabla^\perp \cdot u)|^2 d\mu_g + \mu \int_{\Omega^+} D_t |\nabla^2 (\nabla^\perp \cdot \beta)|^2 d\mu_g \quad (7.10)$$

$$+ \int_{\Gamma} D_t (\gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a P \nabla_F^2 \nabla_f P) \vartheta d\mu_{\gamma} \quad (7.11)$$

$$+ \int_{\Gamma} \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a P \nabla_F^2 \nabla_f P \left(\frac{\vartheta_t}{\vartheta} - h_{NN} \right) \vartheta d\mu_{\gamma}. \quad (7.12)$$

From Lemmas 2.1 and A.3, we get in Ω^+ ,

$$\begin{aligned} & D_t (g^{bd} \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a u_b \nabla_F^2 \nabla_f u_d) \\ &= -2 \nabla_c u_e \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a u^c \nabla_F^2 \nabla_f u^e - 12 \nabla_c u_e \gamma^{ac} \gamma^{ef} \gamma^{AF} \nabla_A^2 \nabla_a u^d \nabla_F^2 \nabla_f u_d \\ &\quad - 2 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f u^b \nabla_A^2 \nabla_a \nabla_b q^+ + 2 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f u^b (\nabla_b u_c - \nabla_c u_b) \nabla_A^2 \nabla_a u^c \\ &\quad + 2 \mu \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f u_d (\beta^c \nabla_c \nabla_{Aa}^3 \beta^d) + 6 \mu \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f u_d (\nabla \beta^c \nabla^2 \nabla_c \beta^d)_{Aa} \\ &\quad + 2 \mu \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f u_d \nabla_{Aa}^3 \beta^c \nabla_c \beta_b - 6 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f u_d \nabla_A^2 u^c \nabla_a \nabla_c u_b \\ &\quad + 6 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f u_d \mu \nabla_A^2 \beta^c \nabla_a \nabla_c \beta_b, \end{aligned}$$

and

$$\begin{aligned} & D_t (g^{bd} \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a \beta_b \nabla_F^2 \nabla_f \beta_d) \\ &= -2 \nabla_c u_e \gamma^{af} \gamma^{AF} \nabla_A^2 \nabla_a \beta^c \nabla_F^2 \nabla_f \beta^e - 6 \nabla_c u_e \gamma^{ac} \gamma^{ef} \gamma^{AF} \nabla_A^2 \nabla_a \beta^d \nabla_F^2 \nabla_f \beta_d \\ &\quad + 2 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f \beta^b (\nabla_b u_c + \nabla_c u_b) \nabla_A^2 \nabla_a \beta^c - 2 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f \beta^b \nabla_c \beta_b \nabla_A^2 \nabla_a u^c \\ &\quad + 2 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f \beta^d \beta^c \nabla_c \nabla_{Aa}^3 u_d + 6 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f \beta^d (\nabla \beta \cdot \nabla^3 u_b)_{Aa} \\ &\quad - 6 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f \beta^d \nabla_A^2 u^c \nabla_a \nabla_c \beta_b + 6 \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f \beta^d \nabla_A^2 \beta^c \nabla_a \nabla_c u_b, \end{aligned}$$

and on the interface Γ ,

$$D_t (\gamma^{af} \gamma^{AF} \nabla_{Aa}^3 P \nabla_{Ff}^3 P) = -6 \nabla_c u_e \gamma^{ac} \gamma^{ef} \gamma^{AF} \nabla_{Aa}^3 P \nabla_{Ff}^3 P + 2 \gamma^{af} \gamma^{AF} \nabla_{Aa}^3 P D_t \nabla_{Ff}^3 P.$$

Thus, we get

$$(7.8) + (7.9) + (7.11) \leq C (\|\nabla u\|_{L^\infty(\Omega^+)} + \|\nabla \beta\|_{L^\infty(\Omega^+)}) E_3(t) \quad (7.13)$$

$$+ C E_3^{1/2}(t) (\|\nabla^2 u\|_{L^\infty(\Omega^+)} + \|\nabla^2 \beta\|_{L^\infty(\Omega^+)}) \cdot (\|\nabla^2 u\|_{L^2(\Omega^+)} + \|\nabla^2 \beta\|_{L^2(\Omega^+)}) \quad (7.14)$$

$$+ 2 \int_{\Gamma} \gamma^{af} \gamma^{AF} \nabla_{Aa}^3 P \left(D_t \nabla_{Ff}^3 P - \frac{1}{\vartheta} N_b \nabla_{Ff}^3 u^b \right) \vartheta d\mu_{\gamma} \quad (7.15)$$

$$- \mu \int_{\Gamma} \gamma^{af} \gamma^{AF} \nabla_{Aa}^3 |\varpi|^2 N_b \nabla_{Ff}^3 u^b d\mu_{\gamma} \quad (7.16)$$

$$+ 2 \int_{\Omega^+} \nabla_b (\gamma^{af} \gamma^{AF}) \nabla_{Ff}^3 u^b \nabla_{Aa}^3 q^+ d\mu_g \quad (7.17)$$

$$+ 2\mu \int_{\Gamma} N_c \gamma^{af} \gamma^{AF} \nabla_F^2 \nabla_f u_d \beta^c \nabla_{Aa}^3 \beta^d d\mu_{\gamma} \quad (7.18)$$

$$- 2\mu \int_{\Omega^+} \nabla_c (\gamma^{af} \gamma^{AF}) \nabla_F^2 \nabla_f u_d \beta^c \nabla_{Aa}^3 \beta^d d\mu_g. \quad (7.19)$$

From Lemma A.7, it follows that

$$|(7.14)| \leq C(K, K_1, \text{Vol } \Omega) \left(E_2(t) E_3^{1/2}(t) + E_2^{1/2}(t) E_3(t) \right).$$

By Gauss' formula, Proposition 3.3 and (3.13), we get

$$\begin{aligned} (7.16) &= -\mu \int_{\Gamma} \gamma^{af} \nabla_f [\gamma^{AF} \nabla_{Aa}^3 |\varpi|^2 N_b \nabla_F^2 u^b - N_a N^c \gamma^{AF} \nabla_{Ac}^3 |\varpi|^2 N_b \nabla_F^2 u^b] d\mu_{\gamma} \\ &\quad + \mu \int_{\Gamma} \gamma^{af} \nabla_f [\gamma^{AF} \nabla_{Aa}^3 |\varpi|^2 N_b] \nabla_F^2 u^b d\mu_{\gamma} \\ &\quad - \mu \int_{\Gamma} \gamma^{af} \nabla_f N_a N^c \gamma^{AF} \nabla_{Ac}^3 |\varpi|^2 N_b \nabla_F^2 u^b d\mu_{\gamma} \\ &\leq \mu C(K, K_1, \text{Vol } \Omega^-) E_0(0) \|\nabla^2 u\|_{L^2(\Gamma)} \\ &\leq \mu C(K, K_1, \text{Vol } \Omega^-, \text{Vol } \Omega^+) E_0(0) (\|\nabla^3 u\|_{L^2(\Omega^+)} + \|\nabla^2 u\|_{L^2(\Omega^+)}) \\ &\leq \mu C(K, K_1, \text{Vol } \Omega) E_0(0) \left(E_3^{1/2}(t) + E_2^{1/2}(t) \right). \end{aligned}$$

From (A.10) and (3.13), it follows

$$\|u\|_{L^\infty(\Gamma)} \leq C(K_1) \sum_{s=0}^2 \|\nabla^s u\|_{L^2(\Omega^+)} \leq C(K_1) \sum_{s=0}^2 E_s^{1/2}(t).$$

From Lemma A.8, it follows, for $\iota_1 \geq 1/K_1$, that

$$\|\beta\|_{L^\infty(\Omega^+)} \leq C \sum_{0 \leq s \leq 2} K_1^{n/2-s} \|\nabla^s \beta\|_{L^2(\Omega^+)} \leq C(K_1) \sum_{s=0}^2 E_s^{1/2}(t). \quad (7.20)$$

Thus, with the help of the Hölder inequality, we have

$$(7.19) \leq CK \|\beta\|_{L^\infty(\Omega^+)} E_3(t) \leq C(K, K_1) \left(\sum_{s=0}^2 E_s^{1/2}(t) \right) E_3(t).$$

From the Hölder inequality, we get

$$(7.17) \leq CK E_3^{1/2}(t) \|\nabla^3 q^+\|_{L^2(\Omega^+)}.$$

We have from (6.26)

$$\nabla \Delta q^+ = \nabla \left(-\nabla_a u^b \nabla_b u^a + \mu \nabla_a \beta^b \nabla_b \beta^a \right) = -2 \nabla \nabla_a u^b \nabla_b u^a + 2 \mu \nabla \nabla_a \beta^b \nabla_b \beta^a.$$

From (7.20), we get for $s \geq 0$

$$\|\nabla^s \beta\|_{L^\infty(\Omega^+)} \leq C \sum_{\ell=0}^2 K_1^{n/2-\ell} \|\nabla^{\ell+s} \beta\|_{L^2(\Omega^+)} \leq C(K_1) \sum_{\ell=0}^2 E_{s+\ell}^{1/2}(t), \quad (7.21)$$

and, similarly,

$$\|\nabla^s u\|_{L^\infty(\Omega^+)} \leq C(K_1) \sum_{\ell=0}^2 E_{s+\ell}^{1/2}(t). \quad (7.22)$$

From Hölder's inequality, (7.21) and (7.22), we get that

$$\begin{aligned} \|\nabla \Delta q^+\|_{L^2(\Omega^+)} &\leq 2 \|\nabla \nabla_a u^b \nabla_b u^a\|_{L^2(\Omega^+)} + 2\mu \|\nabla \nabla_a \beta^b \nabla_b \beta^a\|_{L^2(\Omega^+)} \\ &\leq C \|\nabla u\|_{L^\infty(\Omega^+)} \|\nabla^2 u\|_{L^2(\Omega^+)} + C \|\nabla \beta\|_{L^\infty(\Omega^+)} \|\nabla^2 \beta\|_{L^2(\Omega^+)} \\ &\leq C(K_1) E_2^{1/2}(t) \sum_{\ell=1}^3 E_\ell^{1/2}(t) \\ &\leq C(K_1) \sum_{\ell=1}^2 E_\ell(t) + C(K_1) E_2^{1/2}(t) E_3^{1/2}(t). \end{aligned} \quad (7.23)$$

We have from the assumption (7.3) and Hölder's inequality,

$$\begin{aligned} \|\Delta q^+\|_{L^2(\Omega^+)} &\leq C \|\nabla u\|_{L^2(\Omega^+)} \|\nabla u\|_{L^\infty(\Omega^+)} + C \|\nabla \beta\|_{L^2(\Omega^+)} \|\nabla \beta\|_{L^\infty(\Omega^+)} \\ &\leq C M E_1^{1/2}(t), \end{aligned} \quad (7.24)$$

which is a lower energy term. Thus, by (A.8), (7.23) and (7.24), we obtain

$$\begin{aligned} \|\nabla^2 q^+\|_{L^2(\Gamma)} + \|\nabla^3 q^+\|_{L^2(\Omega^+)} &\leq \|\Pi \nabla^3 q^+\|_{L^2(\Gamma)} + C(K, \text{Vol } \Omega^+) \sum_{s \leq 1} \|\nabla^s \Delta q^+\|_{L^2(\Omega^+)} \\ &\leq \|\Pi \nabla^3 q^+\|_{L^2(\Gamma)} + C(K, K_1, M, \text{Vol } \Omega^+) \sum_{\ell=1}^2 E_\ell(t) \\ &\quad + C(K, K_1, M, \text{Vol } \Omega^+) E_2^{1/2}(t) E_3^{1/2}(t). \end{aligned} \quad (7.25)$$

By (6.25), we have

$$\frac{D_t |\nabla_N P|}{|\nabla_N P|} = -h_{NN} + \frac{\nabla_N D_t P}{\nabla_N P},$$

which implies

$$\ln |\nabla_N P(t)| \leq \ln |\nabla_N P(0)| + (M + L/\varepsilon)t,$$

namely, on Γ

$$|\nabla P(t)| = |\nabla_N P(t)| \leq |\nabla_N P(0)| e^{(M+L/\varepsilon)t} \leq 2 |\nabla_N P(0)|, \quad (7.26)$$

if we take t small enough.

From (7.26), we get

$$\|\Pi \nabla^3 P\|_{L^2(\Gamma)} \leq \|\nabla_N P\|_{L^\infty(\Gamma)}^{1/2} E_3^{1/2}(t) \leq 2 \|\nabla_N P(0)\|_{L^\infty(\Gamma)}^{1/2} E_3^{1/2}(t).$$

Thus, from (7.25) and Proposition 3.3, it follows that

$$\|\nabla^2 q^+\|_{L^2(\Gamma)} + \|\nabla^3 q^+\|_{L^2(\Omega^+)} \leq 2 \|\nabla_N P(0)\|_{L^\infty(\Gamma)}^{1/2} E_3^{1/2}(t) + C(K, K_1, \text{Vol } \Omega^-) E_0(0)$$

$$\begin{aligned}
& + C(K, K_1, M, \text{Vol } \Omega^+) \sum_{\ell=1}^2 E_\ell(t) \\
& + C(K, K_1, M, \text{Vol } \Omega^+) E_2^{1/2}(t) E_3^{1/2}(t). \tag{7.27}
\end{aligned}$$

Then,

$$\begin{aligned}
(7.17) & \leq CK \|\nabla_N P(0)\|_{L^\infty(\Gamma)}^{1/2} E_3(t) + C(K, K_1, \text{Vol } \Omega^-) E_0(0) E_3^{1/2}(t) \\
& + C(K, K_1, M, \text{Vol } \Omega^+) \sum_{\ell=1}^2 E_\ell(t) E_3^{1/2}(t) + C(K, K_1, M, \text{Vol } \Omega^+) E_2^{1/2}(t) E_3(t).
\end{aligned}$$

Now, we turn to the estimates of (7.15). Due to $P = 0$ on Γ implying $\gamma_b^a \nabla_a P = 0$ on Γ , we have from (A.1), by noticing $\vartheta = -1/\nabla_N P$, that

$$-\vartheta^{-1} N_b = \nabla_N P N_b = N^a \nabla_a P N_b = \delta_b^a \nabla_a P - \gamma_b^a \nabla_a P = \nabla_b P. \tag{7.28}$$

By Hölder's inequality and (7.28), we get

$$|(7.15)| \leq C \|\vartheta\|_{L^\infty(\Gamma)}^{1/2} E_3^{1/2}(t) \|\Pi (D_t (\nabla^3 P) + \nabla^3 u \cdot \nabla P)\|_{L^2(\Gamma)}.$$

It follows from (2.11) that

$$D_t \nabla^3 P + \nabla^3 u \cdot \nabla P = -3(\nabla^2 u) \cdot \nabla P + \nabla^3 D_t P. \tag{7.29}$$

We first consider the estimates of the last term in (7.29). By (A.3), (7.32) and (3.13), we have

$$\|\Pi \nabla^3 D_t P\|_{L^2(\Gamma)} \leq \|\bar{\nabla} \theta\|_{L^2(\Gamma)} \|\nabla_N D_t P\|_{L^\infty(\Gamma)} + C \|\theta\|_{L^\infty(\Gamma)} \|\bar{\nabla} \nabla_N D_t P\|_{L^2(\Gamma)}. \tag{7.30}$$

By (A.2) and $P = 0$ on Γ , we have $\Pi \nabla^2 P = \theta \nabla_N P$, i.e., $\theta = -\Pi \nabla^2 P \vartheta$, and then

$$\|\theta\|_{L^2(\Gamma)} = \|\vartheta\|_{L^\infty(\Gamma)}^{1/2} E_2^{1/2}(t) \leq \varepsilon^{-1/2} E_2^{1/2}(t). \tag{7.31}$$

Due to

$$\bar{\nabla}_b \nabla_N P = \theta_b^a \nabla_a P + N^a \nabla_b \nabla_a P - N_b N^d (\theta_d^a \nabla_a P + N^a \nabla_d \nabla_a P),$$

by (7.31), (7.27), (7.26) and (3.13), we have

$$\begin{aligned}
\|\bar{\nabla} \nabla_N P\|_{L^2(\Gamma)} & \leq C \|\theta\|_{L^2(\Gamma)} \|\nabla P\|_{L^\infty(\Gamma)} + C \|\nabla^2 P\|_{L^2(\Gamma)} \\
& \leq C(1/\varepsilon, \|\nabla_N P(0)\|_{L^\infty(\Gamma)}) (E_2^{1/2}(t) + E_3^{1/2}(t)) + C(K, K_1, \text{Vol } \Omega^-) E_0(0) \\
& + C(K, K_1, M, \text{Vol } \Omega^+) \sum_{\ell=1}^2 E_\ell(t) + C(K, K_1, M, \text{Vol } \Omega^+) E_2^{1/2}(t) E_3^{1/2}(t).
\end{aligned}$$

Thus, from (A.3), it yields $(\bar{\nabla} \theta) \nabla_N P = \Pi \nabla^3 P - 3\theta \tilde{\otimes} \bar{\nabla} \nabla_N P$ and then

$$\begin{aligned}
\|\bar{\nabla} \theta\|_{L^2(\Gamma)} & \leq \|\vartheta \Pi \nabla^3 P\|_{L^2(\Gamma)} + \frac{C}{\varepsilon} \|\theta\|_{L^\infty(\Gamma)} \|\bar{\nabla} \nabla_N P\|_{L^2(\Gamma)} \\
& \leq C(1/\varepsilon, \|\nabla_N P(0)\|_{L^\infty(\Gamma)}) (E_2^{1/2}(t) + E_3^{1/2}(t)) + C(K, K_1, \text{Vol } \Omega^-) E_0(0) \\
& + C(K, K_1, M, \text{Vol } \Omega^+) \sum_{\ell=1}^2 E_\ell(t) + C(K, K_1, M, \text{Vol } \Omega^+) E_2^{1/2}(t) E_3^{1/2}(t). \tag{7.32}
\end{aligned}$$

We also have

$$\bar{\nabla}_b \nabla_N D_t P = \theta_b^a \nabla_a D_t P + N^a \nabla_b \nabla_a D_t P - N_b N^d (\theta_d^a \nabla_a D_t P + N^a \nabla_d \nabla_a D_t P),$$

then from (7.30)

$$\begin{aligned} \|\Pi \nabla^3 D_t P\|_{L^2(\Gamma)} &\leq C(1/\varepsilon, K, \|\nabla_N P(0)\|_{L^\infty(\Gamma)}) L(E_2^{1/2}(t) + E_3^{1/2}(t)) \\ &\quad + C(K, K_1, \text{Vol } \Omega^-) L E_0(0) + C(K, K_1, M, \text{Vol } \Omega^+) L \sum_{\ell=1}^2 E_\ell(t) \\ &\quad + C(K, K_1, M, \text{Vol } \Omega^+) L E_2^{1/2}(t) E_3^{1/2}(t) + C \|\nabla^2 D_t P\|_{L^2(\Gamma)}. \end{aligned}$$

By (A.8), it yields

$$\|\nabla^2 D_t q^+\|_{L^2(\Gamma)} \leq \delta \|\Pi \nabla^3 D_t q^+\|_{L^2(\Gamma)} + C(1/\delta, K, \text{Vol } \Omega^+) \sum_{s \leq 1} \|\nabla^s \Delta D_t q^+\|_{L^2(\Omega^+)}. \quad (7.33)$$

From (2.10) and Lemma 2.1, it follows

$$\begin{aligned} \Delta D_t q^+ &= 4g^{ac} \nabla_c u^b \nabla_a \nabla_b q^+ + (\Delta u^e) \nabla_e q^+ + 2\nabla_e u^b \nabla_b u^a \nabla_a u^e \\ &\quad - 2\mu \nabla_b u^a \nabla_a \beta^c \nabla_c \beta^b - 2\mu \nabla_b u^a \beta^c \nabla_a \nabla_c \beta^b + 2\mu \nabla_b \beta^a \beta^e \nabla_e \nabla_a u^b. \end{aligned}$$

By (7.21), (7.25) and Lemma A.8, we get

$$\begin{aligned} \|\Delta D_t q^+\|_{L^2(\Omega^+)} &\leq C \|\nabla u\|_{L^\infty(\Omega^+)} \|\nabla^2 q^+\|_{L^2(\Omega^+)} + C \|\nabla^2 u\|_{L^2(\Omega^+)} \|\nabla q^+\|_{L^\infty(\Omega^+)} \\ &\quad + C \left(\|\nabla u\|_{L^\infty(\Omega^+)}^2 + \|\nabla \beta\|_{L^\infty(\Omega^+)}^2 \right) \|\nabla u\|_{L^2(\Omega^+)} \\ &\quad + C \|\nabla u\|_{L^\infty(\Omega^+)} \|\beta\|_{L^\infty(\Omega^+)} \|\nabla^2 \beta\|_{L^2(\Omega^+)}, \end{aligned}$$

and

$$\begin{aligned} \|\nabla \Delta D_t q^+\|_{L^2(\Omega^+)} &\leq C \|\nabla u\|_{L^\infty(\Omega^+)} \|\nabla^3 q^+\|_{L^2(\Omega^+)} \\ &\quad + C \|\nabla^2 u\|_{L^4(\Omega^+)} \|\nabla^2 q^+\|_{L^4(\Omega^+)} + C \|\nabla^3 u\|_{L^2(\Omega^+)} \|\nabla q^+\|_{L^\infty(\Omega^+)} \\ &\quad + C \left(\|\nabla u\|_{L^\infty(\Omega^+)}^2 + \|\nabla \beta\|_{L^\infty(\Omega^+)}^2 \right) \|\nabla^2 u\|_{L^2(\Omega^+)} \\ &\quad + C \|\nabla u\|_{L^\infty(\Omega^+)} \|\nabla \beta\|_{L^\infty(\Omega^+)} \|\nabla^2 \beta\|_{L^2(\Omega^+)} \\ &\quad + C \|\nabla^2 u\|_{L^4(\Omega^+)} \|\nabla^2 \beta\|_{L^4(\Omega^+)} \|\beta\|_{L^\infty(\Omega^+)} \\ &\quad + C \|\nabla u\|_{L^\infty(\Omega^+)} \|\beta\|_{L^\infty(\Omega^+)} \|\nabla^3 \beta\|_{L^2(\Omega^+)} \\ &\quad + C \|\nabla^3 u\|_{L^2(\Omega^+)} \|\beta\|_{L^\infty(\Omega^+)} \|\nabla \beta\|_{L^\infty(\Omega^+)}. \end{aligned}$$

From Lemma A.7 and (7.22), it follows for $s \leq 2$

$$\begin{aligned} \|\nabla^{s+1} u\|_{L^4(\Omega^+)} &\leq C \|\nabla^s u\|_{L^\infty(\Omega^+)}^{1/2} \left(\sum_{\ell=0}^2 \|\nabla^{s+\ell} u\|_{L^2(\Omega^+)} K_1^{2-\ell} \right)^{1/2} \\ &\leq C(K_1) \sum_{\ell=0}^2 E_{s+\ell}^{1/2}(t). \end{aligned}$$

All the terms with $L^4(\Omega^+)$ norms can be estimated in the same way with the help of (7.21), (7.22), the similar estimates of q^+ and the assumptions. Then, we obtain the bound which is linear about the highest-order derivative or the highest-order energy $E_3^{1/2}(t)$, i.e.,

$$\begin{aligned} \|\nabla^s \Delta D_t q^+\|_{L^2(\Omega^+)} &\leq C(K, K_1, M, M_1, L, 1/\varepsilon, \text{Vol } \Omega^+, E_0(0)) \\ &\quad \cdot \left(1 + \sum_{\ell=0}^2 E_\ell(t) \right) (1 + E_3^{1/2}(t)). \end{aligned} \quad (7.34)$$

Because of

$$D_t q^- = \frac{\mu}{2} D_t |\varpi|^2 = -\mu \varpi^a (\nabla^\perp \Xi)_a + \mu u^c \varpi^a \nabla_c \varpi_a,$$

it follows, from Propositions 3.1 and 4.1 and (A.13), that

$$\begin{aligned} \|\nabla^k D_t q^-\|_{L^2(\Gamma)} &\leq \mu C(K, \text{Vol } \Omega^+) E_0(0) \sum_{s=0}^{k+1} E_s^{1/2}(t) \\ &\quad + \mu C(K, \text{Vol } \Omega^+) E_0(0) \|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)} e^{C(K,L)t}. \end{aligned}$$

Then, from (7.30), (7.33), (7.34) and taking some small δ independent of $E_3(t)$, we obtain

$$\begin{aligned} \|\Pi \nabla^3 D_t q^+\|_{L^2(\Gamma)} &\leq \mu C(K) E_0(0) \|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)} e^{C(K,L)t} \\ &\quad + (1 + \mu) C(K, K_1, M, M_1, L, 1/\varepsilon, \text{Vol } \Omega, E_0(0)) \\ &\quad \cdot \left(1 + \sum_{\ell=0}^2 E_\ell(t)\right) (1 + E_3^{1/2}(t)). \end{aligned} \quad (7.35)$$

Then, we have the similar bound for $\|\Pi \nabla^3 D_t P\|_{L^2(\Gamma)}$.

For (7.29), it only remains to estimate $\|\Pi (\nabla^2 u \cdot \nabla P)\|_{L^2(\Gamma)}$. We get, from (7.6) and Lemma A.10, that

$$\begin{aligned} \|\Pi (\nabla^2 u \cdot \nabla P)\|_{L^2(\Gamma)} &\leq \|\nabla^2 u\|_{L^2(\Gamma)} \|\nabla P\|_{L^\infty(\Gamma)} \\ &\leq C(K, \|\nabla_N P(0)\|_{L^\infty(\Gamma)}, \text{Vol } \Omega^+) L(\text{Vol } \Omega^+)^{1/6} (\|\nabla^3 u\|_{L^2(\Omega^+)} + \|\nabla^2 u\|_{L^2(\Omega^+)}) \\ &\leq C(K, L, \|\nabla_N P(0)\|_{L^\infty(\Gamma)}, \text{Vol } \Omega^+) (E_2^{1/2}(t) + E_3^{1/2}(t)). \end{aligned}$$

Thus, we get

$$\begin{aligned} |(7.15)| &\leq C(K, K_1, M, M_1, L, 1/\varepsilon, \|\nabla_N P(0)\|_{L^\infty(\Gamma)}, \text{Vol } \Omega, E_0(0)) \left(1 + \sum_{s=0}^2 E_s(t)\right) \\ &\quad \cdot (1 + E_3(t)) + \mu C(K, \text{Vol } \Omega^+) E_0(0) \|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)} e^{C(K,L)t} E_3^{1/2}(t). \end{aligned}$$

Therefore, we have obtained

$$\begin{aligned} |(7.8) + (7.9) + (7.11)| &\leq C(K, K_1, M, M_1, L, 1/\varepsilon, \|\nabla_N P(0)\|_{L^\infty(\Gamma)}, \text{Vol } \Omega, E_0(0)) \left(1 + \sum_{s=0}^2 E_s(t)\right) \\ &\quad \cdot (1 + E_3(t)) + \mu C(r, K) E_0(0) \|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)} e^{C(K,L)t} E_3^{1/2}(t). \end{aligned}$$

By a similar argument in Section 6, we get

$$|(7.10)| \leq C(K, K_1, M, \text{Vol } \Omega, 1/\varepsilon) \left(1 + \sum_{s=0}^2 E_s(t)\right) E_3(t).$$

From (6.25), we can easily obtain that (7.12) can be controlled by $C(K, M, L, 1/\varepsilon) E_3(t)$.

Note that there always exists a constant $C > 0$ such that $\|\nabla^\perp \Xi_0\|_{L^2(\Omega^-)}^2 \leq C E_0(0)$ at initial time. Therefore, we obtain

$$\begin{aligned} \frac{d}{dt} E_3(t) &\leq C(K, K_1, M, M_1, L, 1/\varepsilon, \text{Vol } \Omega, E_0(0)) \left(1 + \sum_{s=0}^2 E_s(t)\right) (1 + E_3(t)) \\ &\quad + \mu^2 C(r, K) E_0(0) e^{C(K,L)t}, \end{aligned}$$

which implies, by Grönwall's inequality, that

$$\begin{aligned} E_3(t) \leq & E_3(0) \exp \left(C(K, K_1, M, M_1, L, 1/\varepsilon, \text{Vol } \Omega, E_0(0)) \int_0^t \left(1 + \sum_{s=0}^2 E_s(\tau) \right) d\tau \right) \\ & + \left\{ C(K, K_1, M, M_1, L, 1/\varepsilon, \text{Vol } \Omega, E_0(0)) + \mu^2 e^{C(K, L)t} \right\} \\ & \cdot \int_0^t \left(1 + \sum_{s=0}^2 E_s(\tau) \right) \exp \left(C(K, K_1, M, M_1, L, 1/\varepsilon, \text{Vol } \Omega, E_0(0)) \right. \\ & \quad \left. \cdot \int_\tau^t \left(1 + \sum_{s=0}^2 E_s(s) \right) ds \right) d\tau. \end{aligned}$$

□

8. JUSTIFICATION OF A PRIORI ASSUMPTIONS

Let $\mathcal{K}(t)$ and $\varepsilon(t)$ be the maximum and minimum values, respectively, such that (7.4) and (7.5) hold at time t :

$$\mathcal{K}(t) = \max \left(\|\theta(t, \cdot)\|_{L^\infty(\Gamma)}, 1/\iota_0(t) \right), \quad (8.1)$$

$$\mathcal{E}(t) = \|1/(\nabla_N P(t, \cdot))\|_{L^\infty(\Gamma)} = 1/\varepsilon(t). \quad (8.2)$$

Lemma 8.1. *Assume $\|\theta\|_{L^\infty(W)} + 1/\iota'_0 \leq C\mathcal{K}(0)$. Let $K_1 \geq 1/\iota_1$ be as in Definition A.2, $\mathcal{E}(t)$ as in (8.2). Then there are continuous functions G_j , $j = 1, 2, 3, 4, 5$, such that*

$$\|\nabla u\|_{L^\infty(\Omega)} + \|\nabla \beta\|_{L^\infty(\Omega)} + \|\beta\|_{L^\infty(\Omega)} \leq G_1(K_1, E_0, \dots, E_3), \quad (8.3)$$

$$\|\nabla P\|_{L^\infty(\Omega)} + \|\nabla^2 P\|_{L^\infty(\Gamma)} \leq G_2(K_1, \mathcal{E}, E_0, \dots, E_3, \text{Vol } \Omega), \quad (8.4)$$

$$\|\theta\|_{L^\infty(\Gamma)} \leq G_3(K_1, \mathcal{E}, E_0, \dots, E_3, \text{Vol } \Omega), \quad (8.5)$$

$$\|u\|_{L^\infty(\Gamma)} + \|\nabla D_t P\|_{L^\infty(\Gamma)} \leq G_4(K_1, \mathcal{E}, E_0, \dots, E_3, \text{Vol } \Omega), \quad (8.6)$$

$$\|u\|_{L^\infty(\Gamma)} \leq G_5(K_1, \mathcal{E}, E_0, \dots, E_3, \text{Vol } \Omega). \quad (8.7)$$

Proof. (8.3) follows from (7.22), (7.21) and (7.20). (8.4) follows from Lemmas A.8 and A.6, Lemmas A.9–A.10, and (7.27). Since, from (A.2),

$$|\nabla^2 P| \geq |\Pi \nabla^2 P| = |\nabla_N P| |\theta| \geq \mathcal{E}^{-1} |\theta|, \quad (8.8)$$

(8.5) follows from (8.4). (8.6) follows from Lemma A.6, (7.33), (7.34) and (7.35). (8.7) follows from Lemmas A.6 and A.10. □

Lemma 8.2. *Assume $\|\theta\|_{L^\infty(W)} + 1/\iota'_0 \leq C\mathcal{K}(0)$. Let $K_1 \geq 1/\iota_1$ and ε_1 be as in Definition A.2. Then*

$$\left| \frac{d}{dt} E_3 \right| \leq C(K_1, \mathcal{E}, E_0, \dots, E_3, \text{Vol } \Omega) \sum_{s=0}^3 E_s, \quad (8.9)$$

and

$$\left| \frac{d}{dt} \mathcal{E} \right| \leq C(K_1, \mathcal{E}, E_0, \dots, E_3, \text{Vol } \Omega). \quad (8.10)$$

Proof. (8.9) is a consequence of Lemma 8.1 and the estimates in the proof of Theorems 7.1. (8.10) follows from (8.6) and

$$\left| \frac{d}{dt} \left\| \frac{1}{-\nabla_N P(t, \cdot)} \right\|_{L^\infty(\Gamma)} \right| \leq C \left\| \frac{1}{-\nabla_N P(t, \cdot)} \right\|_{L^\infty(\Gamma)}^2 \|\nabla_N D_t P(t, \cdot)\|_{L^\infty(\Gamma)}.$$

□

As a consequence of Lemma 8.2, we have the following:

Lemma 8.3. *Assume $\|\theta\|_{L^\infty(W)} + 1/\iota'_0 \leq CK(0)$. There exists a continuous function $\mathcal{T} > 0$ depending on $K_1, \mathcal{E}(0), E_0(0), \dots, E_3(0), \text{Vol } \Omega$ such that for*

$$0 \leq t \leq \mathcal{T}(K_1, \mathcal{E}(0), E_0(0), \dots, E_3(0), \text{Vol } \Omega),$$

the following statements hold: We have

$$E_s(t) \leq 2E_s(0), \quad 0 \leq s \leq 3; \quad \mathcal{E}(t) \leq 2\mathcal{E}(0). \quad (8.11)$$

Furthermore,

$$\frac{g_{ab}(0, y)Y^a Y^b}{2} \leq g_{ab}(t, y)Y^a Y^b \leq 2g_{ab}(0, y)Y^a Y^b, \quad (8.12)$$

and with ε_1 as in Definition A.2,

$$|\mathbf{n}(x(t, \bar{y})) - \mathbf{n}(x(0, \bar{y}))| \leq \frac{\varepsilon_1}{16}, \quad \bar{y} \in \Gamma, \quad (8.13)$$

$$|x(t, y) - x(0, y)| \leq \frac{\iota_1}{16}, \quad y \in \Omega, \quad (8.14)$$

$$\left| \frac{\partial x(t, \bar{y})}{\partial y} - \frac{\partial x(0, \bar{y})}{\partial y} \right| \leq \frac{\varepsilon_1}{16}, \quad \bar{y} \in \Gamma. \quad (8.15)$$

Proof. Since the proof is similar to [9, Lemma 6.3], we omit details. $\square$

Now we use (8.12)–(8.15) to pick a K_1 , i.e., ι_1 , which depends only on its value at $t = 0$,

$$\iota_1(t) \geq \iota_1(0)/2.$$

Lemma 8.4. *Assume $\|\theta\|_{L^\infty(W)} + 1/\iota'_0 \leq CK(0)$. Let $\mathcal{T}$ be as in Lemma 8.2. Pick $\iota_1 > 0$ such that*

$$|\mathbf{n}(x(0, y_1)) - \mathbf{n}(x(0, y_2))| \leq \frac{\varepsilon_1}{2}, \quad \text{whenever } |x(0, y_1) - x(0, y_2)| \leq 2\iota_1. \quad (8.16)$$

Then if $t \leq \mathcal{T}$, we have

$$|\mathbf{n}(x(t, y_1)) - \mathbf{n}(x(t, y_2))| \leq \varepsilon_1, \quad \text{whenever } |x(t, y_1) - x(t, y_2)| \leq 2\iota_1. \quad (8.17)$$

Proof. (8.17) follows from (8.16), (8.13) and (8.14) in view of triangle inequalities. $\square$

Finally, Lemma 8.4 allows us to pick a K_1 depending only on initial conditions, while Lemma 8.3 gives us $\mathcal{T} > 0$ that depends only on the initial conditions and K_1 such that, by Lemma 8.4, $1/\iota_1 \leq K_1$ for $t \leq \mathcal{T}$. Thus, we immediately obtain Theorem 1.1.

APPENDIX A. PRELIMINARIES AND SOME ESTIMATES

Let N^a denote the unit normal to Γ , $g_{ab}N^aN^b = 1$, $g_{ab}N^aT^b = 0$ if $T \in T(\Gamma)$, and let $N_a = g_{ab}N^b$ denote the unit conormal, $g^{ab}N_aN_b = 1$. The induced metric γ on the tangent space to the boundary $T(\Gamma)$ and $T(W)$ extended to be 0 on the orthogonal complement in $T(\Omega)$ is then given by

$$\gamma_{ab} = g_{ab} - N_aN_b, \quad \gamma^{ab} = g^{ab} - N^aN^b.$$

We can extend the normal to a vector field $\mathbf{n}$ defined and regular everywhere in the interior. From [4, Lemma 3.10], it holds $|\nabla \mathbf{n}| \leq 2\|\theta\|_{L^\infty(\Gamma \cup W)}$. One can see [4] for more details.

The orthogonal projection of an (r, s) tensor S to the boundary is given by

$$(\Pi S)_{b_1 \dots b_s}^{a_1 \dots a_r} = \gamma_{c_1}^{a_1} \dots \gamma_{c_r}^{a_r} \gamma_{b_1}^{d_1} \dots \gamma_{b_s}^{d_s} S_{d_1 \dots d_s}^{c_1 \dots c_r},$$

where

$$\gamma_a^c = \delta_a^c - N_a N^c. \quad (\text{A.1})$$

Covariant differentiation on the boundary $\bar{\nabla}$ is given by

$$\bar{\nabla} S = \Pi \nabla S.$$

The second fundamental form of the boundary is given by

$$\theta_{ab} = (\Pi \nabla N)_{ab} = \gamma_a^c \nabla_c N_b.$$

Let us now recall some properties of the projection. Since $g^{ab} = \gamma^{ab} + N^a N^b$, we have

$$\Pi(S \cdot R) = \Pi(S) \cdot \Pi(R) + \Pi(S \cdot N) \tilde{\otimes} \Pi(N \cdot R),$$

where $S \tilde{\otimes} R$ denotes some partial symmetrization of the tensor product $S \otimes R$, i.e., a sum over some subset of the permutations of the indices divided by the number of permutations in that subset. Similarly, we let $S \tilde{\cdot} R$ denote a partial symmetrization of the dot product $S \cdot R$. Now we recall some identities:

$$\Pi \nabla^2 q = \bar{\nabla}^2 q + \theta \nabla_N q, \quad (\text{A.2})$$

$$\Pi \nabla^3 q = \bar{\nabla}^3 q - 2\theta \tilde{\otimes} (\theta \cdot \bar{\nabla} q) + (\bar{\nabla} \theta) \nabla_N q + 3\theta \tilde{\otimes} \bar{\nabla} \nabla_N q. \quad (\text{A.3})$$

Definition A.1. Let $\mathbf{n}(\bar{x})$ be the outward unit normal to Γ_t at $\bar{x} \in \Gamma_t$. Let $\text{dist}(x_1, x_2) = |x_1 - x_2|$ denote the Euclidean distance in $\mathbb{R}^n$, and for $\bar{x}_1, \bar{x}_2 \in \Gamma_t$, let $\text{dist}_{\Gamma_t}(\bar{x}_1, \bar{x}_2)$ denote the geodesic distance on the boundary.

Definition A.2. Let $0 < \varepsilon_1 < 2$ be a fixed number, and let $\iota_1 = \iota_1(\varepsilon_1)$ the largest number such that

$$|\mathbf{n}(\bar{x}_1) - \mathbf{n}(\bar{x}_2)| \leq \varepsilon_1 \quad \text{whenever} \quad |\bar{x}_1 - \bar{x}_2| \leq \iota_1, \quad \bar{x}_1, \bar{x}_2 \in \Gamma_t.$$

Lemma A.3 (cf. [4, Lemma 3.9]). *Let N be the unit normal to Γ , and let $h_{ab} = \frac{1}{2} D_t g_{ab}$. On $[0, T] \times \Gamma$, we have*

$$D_t N_a = h_{NN} N_a, \quad D_t N^c = -2h_d^c N^d + h_{NN} N^c, \quad D_t \gamma^{ab} = -2\gamma^{ac} h_{cd} \gamma^{db}, \quad (\text{A.4})$$

where $h_{NN} = h_{ab} N^a N^b$. The volume element on Γ satisfies

$$D_t d\mu_\gamma = (\text{tr } h - h_{NN}) d\mu_\gamma. \quad (\text{A.5})$$

Lemma A.4 (cf. [4, Lemma 5.5]). *Let $w_a = w_{Aa} = \nabla_A^r f_a$, $\nabla_A^r = \nabla_{a_1} \cdots \nabla_{a_r}$, f be a $(0, 1)$ tensor, and $[\nabla_a, \nabla_b] = 0$. Let $\text{div } w = \nabla_a w^a = \nabla^r \text{div } f$, and let $(\nabla \times w)_{ab} = \nabla_a w_b - \nabla_b w_a = \nabla^r (\nabla \times f)_{ab}$. Then,*

$$|\nabla w|^2 \leq C(g^{ab} \gamma^{cd} \gamma^{AB} \nabla_c w_{Aa} \nabla_d w_{Bb} + |\text{div } w|^2 + |\nabla \times w|^2), \quad (\text{A.6})$$

$$\int_\Omega |\nabla w|^2 d\mu_g \leq C \int_\Omega (N^a N^f g^{cd} \gamma^{AF} \nabla_c w_{Aa} \nabla_d w_{Ff} + |\text{div } w|^2 + |\nabla \times w|^2 + K^2 |w|^2) d\mu_g. \quad (\text{A.7})$$

Lemma A.5 (cf. [4, Proposition 5.8]). *Let ι_0 and ι_1 be as in (1.7) and Definition A.2, and suppose that $|\theta| + 1/\iota_0 \leq K$ and $1/\iota_1 \leq K_1$. Then with $\tilde{K} = \min(K, K_1)$ we have, for any $r \geq 2$ and $\delta > 0$,*

$$\|\nabla^{r-1} q\|_{L^2(\Gamma)} + \|\nabla^r q\|_{L^2(\Omega)} \leq \delta \|\Pi \nabla^r q\|_{L^2(\Gamma)} + C(1/\delta, K, \text{Vol } \Omega) \sum_{s \leq r-2} \|\nabla^s \Delta q\|_{L^2(\Omega)}. \quad (\text{A.8})$$

Lemma A.6 ([4, Lemma A.2]). *Suppose that for $\iota_1 \geq 1/K_1$*

$$|\mathbf{n}(\bar{x}_1) - \mathbf{n}(\bar{x}_2)| \leq \varepsilon_1, \quad \text{whenever } |\bar{x}_1 - \bar{x}_2| \leq \iota_1, \quad \bar{x}_1, \bar{x}_2 \in \Gamma_t,$$

and

$$C_0^{-1} \gamma_{ab}^0(y) Z^a Z^b \leq \gamma_{ab}(t, y) Z^a Z^b \leq C_0 \gamma_{ab}^0(y) Z^a Z^b, \quad \text{if } Z \in T(\Omega^+),$$

where $\gamma_{ab}^0(y) = \gamma_{ab}(0, y)$. Then if α is a $(0, r)$ tensor,

$$\|\alpha\|_{L^{(n-1)p/(n-1-kp)}(\Gamma)} \leq C(K_1) \sum_{\ell=0}^k \|\nabla^\ell \alpha\|_{L^p(\Gamma)}, \quad 1 \leq p < \frac{n-1}{k}, \quad (\text{A.9})$$

$$\|\alpha\|_{L^\infty(\Gamma)} \leq \delta \|\nabla^k \alpha\|_{L^p(\Gamma)} + C_\delta(K_1) \sum_{\ell=0}^{k-1} \|\nabla^\ell \alpha\|_{L^p(\Gamma)}, \quad k > \frac{n-1}{p}, \quad (\text{A.10})$$

for any $\delta > 0$.

Lemma A.7 ([4, Lemma A.3]). *If α is a $(0, r)$ tensor, then with $a = k/m$ and a constant C that only depends on m and n , with notation as in Lemma A.6, we have*

$$\sum_{j=0}^k \|\nabla^j \alpha\|_{L^s(\Omega)} \leq C \|\alpha\|_{L^q(\Omega)}^{1-a} \left(\sum_{i=0}^m \|\nabla^i \alpha\|_{L^p(\Omega)} K_1^{m-i} \right)^a,$$

if

$$\frac{m}{s} = \frac{k}{p} + \frac{m-k}{q}, \quad 2 \leq p \leq s \leq q \leq \infty.$$

Lemma A.8 ([4, Lemma A.4]). *Suppose that $\iota_1 \geq 1/K_1$ and α is a $(0, r)$ tensor. Then*

$$\|\alpha\|_{L^{np/(n-kp)}(\Omega)} \leq C \sum_{\ell=0}^k K_1^{k-\ell} \|\nabla^\ell \alpha\|_{L^p(\Omega)}, \quad 1 \leq p < \frac{n}{k}, \quad (\text{A.11})$$

$$\|\alpha\|_{L^\infty(\Omega)} \leq C \sum_{\ell=0}^k K_1^{n/p-\ell} \|\nabla^\ell \alpha\|_{L^p(\Omega)}, \quad k > \frac{n}{p}. \quad (\text{A.12})$$

Lemma A.9 ([4, Lemma A.5]). *Suppose that $q = 0$ on Γ . Then*

$$\|q\|_{L^2(\Omega)} \leq C(\text{Vol } \Omega)^{1/n} \|\nabla q\|_{L^2(\Omega)}, \quad \|\nabla q\|_{L^2(\Omega)} \leq C(\text{Vol } \Omega)^{1/2n} \|\Delta q\|_{L^2(\Omega)}.$$

Lemma A.10 ([4, Lemma A.7]). *Let α be a $(0, r)$ tensor. Assume that $\text{Vol } \Omega \leq V$ and $\|\theta\|_{L^\infty(\Gamma)} + 1/\iota_0 \leq K$, then there is a $C = C(K, V, r, n)$ such that*

$$\|\alpha\|_{L^{(n-1)p/(n-p)}(\Gamma)} \leq C \|\nabla \alpha\|_{L^p(\Omega)} + C \|\alpha\|_{L^p(\Omega)}, \quad 1 \leq p < n. \quad (\text{A.13})$$

ACKNOWLEDGMENTS

Hao's research was supported in part by the National Natural Science Foundation of China under grants 11171327 and 11671384.

REFERENCES

- [1] D. M. Ambrose and N. Masmoudi. The zero surface tension limit of two-dimensional water waves. *Comm. Pure Appl. Math.*, 58(10):1287–1315, 2005.
- [2] J. T. Beale, T. Y. Hou, and J. S. Lowengrub. Growth rates for the linearized motion of fluid interfaces away from equilibrium. *Comm. Pure Appl. Math.*, 46(9):1269–1301, 1993.
- [3] G.-Q. Chen and Y.-G. Wang. Existence and stability of compressible current-vortex sheets in three-dimensional magnetohydrodynamics. *Arch. Ration. Mech. Anal.*, 187(3):369–408, 2008.
- [4] D. Christodoulou and H. Lindblad. On the motion of the free surface of a liquid. *Comm. Pure Appl. Math.*, 53(12):1536–1602, 2000.

- [5] D. Coutand and S. Shkoller. Well-posedness of the free-surface incompressible Euler equations with or without surface tension. *J. Amer. Math. Soc.*, 20(3):829–930, 2007.
- [6] D. G. Ebin. The equations of motion of a perfect fluid with free boundary are not well posed. *Comm. Partial Differential Equations*, 12(10):1175–1201, 1987.
- [7] J. P. Freidberg. *Ideal MHD*. Cambridge University Press, June 2014.
- [8] A. Friedman and Y. Liu. A free boundary problem arising in magnetohydrodynamic system. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 22(3):375–448, 1995.
- [9] C. Hao and T. Luo. A priori estimates for free boundary problem of incompressible inviscid magnetohydrodynamic flows. *Arch. Ration. Mech. Anal.*, 212(3):805–847, 2014.
- [10] D. Lee. Initial value problem for the free boundary magnetohydrodynamics with zero magnetic boundary condition. arXiv:1409.4392, 2014.
- [11] D. Lee. Zero kinetic viscosity- magnetic diffusivity limit of free boundary magnetohydrodynamics. arXiv:1409.5384, 2014.
- [12] H. Lindblad. Well-posedness for the linearized motion of an incompressible liquid with free surface boundary. *Comm. Pure Appl. Math.*, 56(2):153–197, 2003.
- [13] H. Lindblad. Well-posedness for the motion of an incompressible liquid with free surface boundary. *Ann. of Math. (2)*, 162(1):109–194, 2005.
- [14] H. Lindblad and K. H. Nordgren. A priori estimates for the motion of a self-gravitating incompressible liquid with free surface boundary. *J. Hyperbolic Differ. Equ.*, 6(2):407–432, 2009.
- [15] A. Morando, Y. Trakhinin, and P. Trebeschi. Well-posedness of the linearized plasma-vacuum interface problem in ideal incompressible MHD. *Quart. Appl. Math.*, 72(3):549–587, 2014.
- [16] M. Padula and V. A. Solonnikov. On the free boundary problem of magnetohydrodynamics. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 385(Kraevye Zadachi Matematicheskoi Fizikii Smezhnye Voprosy Teorii Funktsii. 41):135–186, 2010. translation in *J. Math. Sci. (N. Y.)* **178**(3), 313–344 (2011).
- [17] D. D. Schnack. *Lectures in magnetohydrodynamics*, volume 780 of *Lecture Notes in Physics*. Springer-Verlag, Berlin, 2009.
- [18] P. Secchi and Y. Trakhinin. Well-posedness of the linearized plasma-vacuum interface problem. *Interfaces Free Bound.*, 15(3):323–357, 2013.
- [19] P. Secchi and Y. Trakhinin. Well-posedness of the plasma-vacuum interface problem. *Nonlinearity*, 27(1):105–169, 2014.
- [20] J. Shatah and C. Zeng. Geometry and a priori estimates for free boundary problems of the Euler equation. *Comm. Pure Appl. Math.*, 61(5):698–744, 2008.
- [21] Y. Sun, W. Wang, and Z. Zhang. Nonlinear stability of current-vortex sheet to the incompressible mhd equations, 2015. arXiv:1510.02228v1.
- [22] Y. Trakhinin. The existence of current-vortex sheets in ideal compressible magnetohydrodynamics. *Arch. Ration. Mech. Anal.*, 191(2):245–310, 2009.
- [23] Y. Trakhinin. On the well-posedness of a linearized plasma-vacuum interface problem in ideal compressible MHD. *J. Differential Equations*, 249(10):2577–2599, 2010.
- [24] S. Wu. Well-posedness in Sobolev spaces of the full water wave problem in 2-D. *Invent. Math.*, 130(1):39–72, 1997.
- [25] S. Wu. Well-posedness in Sobolev spaces of the full water wave problem in 3-D. *J. Amer. Math. Soc.*, 12(2):445–495, 1999.
- [26] P. Zhang and Z. Zhang. On the free boundary problem of three-dimensional incompressible Euler equations. *Comm. Pure Appl. Math.*, 61(7):877–940, 2008.

INSTITUTE OF MATHEMATICS, ACADEMY OF MATHEMATICS & SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA

E-mail address: hcc@amss.ac.cn